

STANDARD SINGLE VALUED NEUTROSOPHIC METRIC SPACES WITH APPLICATION

O. BARKAT¹, S. MILLES^{1*}, A. LATRECHE², §

ABSTRACT. In recent years, J. R. Kider and Z.A. Hussain have introduced the notion of standard fuzzy metric space and some basic properties were proved. In this paper, we generalize this notion to the setting of single valued neutrosophic sets. Furthermore, some interesting properties related to this notion are studied such as the continuity property of the mappings defined on standard single valued neutrosophic metric spaces. As an application, we establish the Edelstein fixed point theorem for standard single valued neutrosophic metric spaces.

Keywords: single valued neutrosophic set; metric space; continuous mapping; fixed point property.

AMS Subject Classification: 03B52, 54E35, 47H10.

1. INTRODUCTION

A metric space is a particular type of topological space where the topology is defined by a distance function. It is one of the important basic areas of research for the mathematicians. In 1906, Maurice Fréchet in his doctoral thesis [9] introduced the concept of a metric space. However, the name is due to Felix Hausdorff. Later on, many researchers developed this concept either by introducing different contractions in different fields or by extending number of variables in it. Metric spaces have applications in many mathematical branches, as well as in topology and its analysis approaches (e.g. [2]), they appear also in differential geometry (e.g. [31]), linear algebra (e.g. [12]), geographic information systems (e.g. [8]), computer science (e.g. [19]) and in coding theory (e.g. [7]).

Recently, Smarandache [23] generalized the concepts of intuitionistic fuzzy sets and other types of sets to the notion of neutrosophic sets. He introduce this concept to know the correct way of dealing with imprecise and indeterminate data. Neutrosophic sets are characterized by three independent components such that truth membership function (T),

¹ Department of Mathematics and Informatics, University Center of Barika, Algeria.
e-mail: omarbarkat@cu-barika.dz; ORCID: <https://orcid.org/0000-0002-1732-4824>.
e-mail: soheybmilles@cu-barika.dz; ORCID: <https://orcid.org/0000-0001-8801-4607>.

* Corresponding author.

² Department of Mathematics, University of Skikda, Algeria.
e-mail: a.latreche@univ-skikda.dz; ORCID: <https://orcid.org/0000-0003-1424-7197>.

§ Manuscript received: August 09, 2021; accepted: November 16, 2021.

TWMS Journal of Applied and Engineering Mathematics, Vol.13, No.3 © Işık University, Department of Mathematics, 2023; all rights reserved.

indeterminacy membership function (I) and falsity membership function (F). Many researchers like Abdel-Basset et al. , Broumi et al. [3,4] and others have studied and applied in various fields the neutrosophic sets and its different extensions such as medical diagnosis (e.g. [34,36]), management (e.g. [1]), decision making problems (e.g. [10,35]), image processing (e.g. [5]), educational problem (e.g. [18]), conflict resolution (e.g. [20]) and in social problems (e.g. [21]). In particular, Wang et al. [32] introduced the notion of single valued neutrosophic set as an instance of neutrosophic set which can be used in real scientific and engineering applications. Since its introduction, it has become omnipresent, where the studies whether theoretical or applied on the single valued neutrosophic set have been advancing at an accelerated pace. For more details on single valued neutrosophic set and background, the readers are referred to [13,17] and more others.

From the idea of neutrosophic sets, Kirişci and Şimşek [15] defined the neutrosophic metric spaces (NMS). They investigated some of their properties and proved that Baire category theorem and uniform convergence theorem. Thereafter, Kirişci et al. [16] introduced the neutrosophic contractive mapping and gave a fixed point results in complete neutrosophic metric spaces. Şimşek and Kirişci [26] discussed some fixed point results in (NMS), where they defined new infinite products by continuous t -norms and proved the Banach contraction theorem for (NMS) by these products. Moreover, there are many researchers have created other types of metric spaces in the context of the neutrosophic theory, where the latter also still knows many developments such as Şahin and Kargin [27] introduced the notion of neutrosophic triplet v -generalized metric space (NTVGM) in which that the concept of neutrosophic triplet (NT) is a new theory in neutrosophy, and the v -generalized metric is a specific form of the classical metrics. In this study, they proved that (NTVGM) is different from the classical metric and neutrosophic triplet metric (NTM). In [30], Taş et al. defined the neutrosophic valued metric spaces and the neutrosophic valued g -metric spaces. Furthermore, they determined a mathematical model for clustering the neutrosophic big data sets with the use of g -metric. In this regard, we find that other authors have adopted the same approach, such as Şahin et al. [28,29].

Motivated by the recent relevant developments to this context, in the present paper, we introduce the notion of standard single valued neutrosophic metric space as a generalization of standard fuzzy metric space introduced by Kider and Hussain [14] and many fundamental properties related to this concept are investigated, especially, the continuity property because of its frequent use in many applications such as in fixed point theory. As an application, we extend the Edelstein fixed point theorem [6] to the standard single valued neutrosophic metric spaces. It is worth noting here that the adoption of the new definition of neutrosophic metric space which we called standard single valued neutrosophic metric space, allowed us to obtain many results and demonstrated them smoothly to a large extent. Specifically, during the generalization from the fuzzy case.

This paper is structured as follows. In Section 2, we recall the necessary basic notions and properties of standard fuzzy metric space and single valued neutrosophic sets with some related concepts that will be needed throughout this paper. In Section 3, the notion of standard fuzzy metric space is introduced and some fundamental properties related to this concept are studied. Subsequently, several interesting properties of single valued neutrosophic continuous mappings in standard single valued neutrosophic metric space are investigated. As an application, Section 4 is devoted to explaining a generalization of Edelstein's fixed point theorem in a standard single value neutrosophical metric space. We end with some conclusions and discuss future work in Section 5.

2. PRELIMINARIES

This section contains the basic definitions and properties of single valued neutrosophic sets and some related notions that will be needed throughout this paper.

2.1. Standard fuzzy metric space. The standard fuzzy metric space has been mentioned as an example of fuzzy metric in a particular case by George and Veeramani [11] and its definition was introduced with studied extensively by Kider and Hussain [14].

Definition 2.1. [14] *A triple $(X, M, *)$ is said to be standard fuzzy metric space if X is an arbitrary set, $*$ is a continuous t -norm and M is a continuous fuzzy set on X^2 satisfying the following conditions:*

- (i) $\mu_M(x, y) > 0$ for all $x, y \in X$;
- (ii) $\mu_M(x, y) = 1$ if and only if $x = y$;
- (iii) $\mu_M(x, y) = \mu_M(y, x)$ for all $x, y \in X$;
- (iv) $\mu_M(x, z) \geq \mu_M(x, y) * \mu_M(y, z)$ for all $x, y, z \in X$.

Example 2.1. Let $X = \mathbb{R}$. Define the t -norm $a * b = ab$, for all $a, b \in [0, 1]$ and let M be fuzzy set on X^2 defined as: $\mu(x, y) = \exp^{-|x-y|}$. It is clear that all the conditions from (i) to (iv) realized. Therefore, $(X, M, *)$ is a standard fuzzy metric space.

2.2. Single valued neutrosophic sets. In 1998, Smarandache [23] defined the concept of a neutrosophic set as a generalization of Atanassov's intuitionistic fuzzy set. Also, he introduced neutrosophic logic, neutrosophic set and its applications in [24, 25]. In particular, Wang et al. [32] introduced the notion of a single valued neutrosophic set.

Definition 2.2. [24] *Let X be a nonempty set. A neutrosophic set (NS, for short) A on X is an object of the form $A = \{\langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle \mid x \in X\}$ characterized by a membership function $\mu_A : X \rightarrow]-0, 1^+[$ and an indeterminacy function $\sigma_A : X \rightarrow]-0, 1^+[$ and a non-membership function $\nu_A : X \rightarrow]-0, 1^+[$ which satisfy the condition:*

$$-0 \leq \mu_A(x) + \sigma_A(x) + \nu_A(x) \leq 3^+, \text{ for any } x \in X.$$

Next, we show the notion of single valued neutrosophic set as an instance of neutrosophic set which can be used in real scientific and engineering applications.

Definition 2.3. [32] *Let X be a nonempty set. A single valued neutrosophic set (SVNS, for short) A on X is an object of the form $A = \{\langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle \mid x \in X\}$ characterized by a truth-membership function $\mu_A : X \rightarrow [0, 1]$, an indeterminacy-membership function $\sigma_A : X \rightarrow [0, 1]$ and a falsity-membership function $\nu_A : X \rightarrow [0, 1]$. A continuous single valued neutrosophic set can be defined as $A = \int_X \langle x, \mu_A(x), \sigma_A(x), \nu_A(x) \rangle \mid x \in X$.*

The class of single valued neutrosophic sets on X is denoted by $SVN(X)$.

For any two SVNSs A and B on a set X , several operations are defined (see, e.g., [32, 33]). Here we will present only those which are related to the present paper.

- (i) $A \subseteq B$ if $\mu_A(x) \leq \mu_B(x)$ and $\sigma_A(x) \leq \sigma_B(x)$ and $\nu_A(x) \geq \nu_B(x)$, for all $x \in X$,
- (ii) $A = B$ if $\mu_A(x) = \mu_B(x)$ and $\sigma_A(x) = \sigma_B(x)$ and $\nu_A(x) = \nu_B(x)$, for all $x \in X$,
- (iii) $A \cap B = \{\langle x, \mu_A(x) \wedge \mu_B(x), \sigma_A(x) \wedge \sigma_B(x), \nu_A(x) \vee \nu_B(x) \rangle \mid x \in X\}$,
- (iv) $A \cup B = \{\langle x, \mu_A(x) \vee \mu_B(x), \sigma_A(x) \vee \sigma_B(x), \nu_A(x) \wedge \nu_B(x) \rangle \mid x \in X\}$,
- (v) $\bar{A} = \{\langle x, 1 - \nu_A(x), 1 - \sigma_A(x), 1 - \mu_A(x) \rangle \mid x \in X\}$,
- (vi) $[A] = \{\langle x, \mu_A(x), \sigma_A(x), 1 - \mu_A(x) \rangle \mid x \in X\}$,
- (vii) $\langle A \rangle = \{\langle x, 1 - \nu_A(x), \sigma_A(x), \nu_A(x) \rangle \mid x \in X\}$.

3. STANDARD SINGLE VALUED NEUTROSOPHIC METRIC SPACE

In this section, we generalize the notion of standard fuzzy metric space introduced by Kider and Hussain [14] to the setting of single valued neutrosophic sets. Also, the main properties related to standard single valued neutrosophic metric space are studied.

3.1. Definitions.

Definition 3.1. A quintuple $(X, M, *, \triangleleft, \diamond)$ is said to be a standard single valued neutrosophic metric space if X is an arbitrary set, $*$, \triangleleft are a continuous t -norms, \diamond is a t -conorm and M is a continuous single valued neutrosophic set on X^2 satisfying the following conditions:

- (i) $\mu_M(x, y) > 0$, $\sigma_M(x, y) > 0$ and $\nu_M(x, y) < 1$ for all $x, y \in X$;
- (ii) $\mu_M(x, y) = 1$, $\sigma_M(x, y) = 1$ and $\nu_M(x, y) = 0$ if and only if $x = y$;
- (iii) $\mu_M(x, y) = \mu_M(y, x)$, $\sigma_M(x, y) = \sigma_M(y, x)$ and $\nu_M(x, y) = \nu_M(y, x)$ for all $x, y \in X$;
- (iv) $\mu_M(x, z) \geq \mu_M(x, y) * \mu_M(y, z)$, $\sigma_M(x, z) \geq \sigma_M(x, y) \triangleleft \sigma_M(y, z)$ and $\nu_M(x, z) \leq \nu_M(x, y) \diamond \nu_M(y, z)$ for all $x, y \in X$.

Here, $\mu_M(x, y)$, $\sigma_M(x, y)$ and $\nu_M(x, y)$ denotes the degree of nearness, the degree of neutrality, and the degree of non-nearness between x and y , respectively.

Remark 3.1. If the set X given in the previous definition is a metric space with an ordinary distance d , then $(X, M, *, \triangleleft, \diamond)$ is called a standard single valued neutrosophic metric space induced by (X, d) , and the single valued neutrosophic distance M is called a standard if it is induced by the ordinary distance d .

Example 3.1. Let (X, d) be an ordinary metric space. Define the t -norms $*$ and \triangleleft by: $x * y = \min\{x, y\}$, $x \triangleleft y = \min\{x, y\}$, and the t -conorm \diamond by: $x \diamond y = \max\{x, y\}$, for all $x, y \in [0, 1]$. Define the single valued neutrosophic set M on X^2 as:

$$\mu_M(x, y) = \frac{1}{1+d(x, y)}, \quad \sigma_M(x, y) = 1 + d(x, y), \quad \nu_M(x, y) = \frac{d(x, y)}{1+d(x, y)}.$$

Then, $(X, M, *, \triangleleft, \diamond)$ is a standard single valued neutrosophic metric space.

Remark 3.2. Let $*$ be a t -norm, \diamond be a t -conorm and $r_1, r_2 \in [0, 1]$. If $r_1 > r_2$, then there exist $r_3, r_4 \in]0, 1[$ such that $r_1 * r_3 \geq r_2$ and $r_1 \geq r_2 \diamond r_4$. Moreover, for any $r_5 \in]0, 1[$, there exist $r_6, r_7 \in]0, 1[$ such that $r_6 * r_7 \geq r_5$ and $r_7 \diamond r_7 \leq r_5$.

Remark 3.3. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. Then, μ_M, σ_M are a monotone mappings, and ν_M is an antitone mapping.

Next, we introduce the standard single valued neutrosophic distance between an element and a subset of X and the standard single valued neutrosophic distance between two subsets of X .

Definition 3.2. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. For $x \in X$ and A, B are two subsets of X . Then

- (i) the standard single valued neutrosophic distance between x and A is defined as

$$\mu_M(x, A) = \inf\{\mu_M(x, y) \mid y \in A\}, \quad \sigma_M(x, A) = \inf\{\sigma_M(x, y) \mid y \in A\}$$

$$\text{and } \nu_M(x, A) = \sup\{\nu_M(x, y) \mid y \in A\};$$

- (ii) the standard single valued neutrosophic distance between A and B is defined as

$$\mu_M(A, B) = \inf\{\mu_M(x, y) \mid x \in A, y \in B\}, \quad \sigma_M(A, B) = \inf\{\sigma_M(x, y) \mid x \in A, y \in B\}$$

$$\text{and } \nu_M(A, B) = \sup\{\nu_M(x, y) \mid x \in A, y \in B\}.$$

Definition 3.3. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. For $x \in X$ and $r \in]0, 1[$, the open ball $\mathcal{B}(x, r)$ with radius r and center x is defined by

$$\mathcal{B}(x, r) = \{y \in X \mid \mu_M(x, y) > 1 - r, \sigma_M(x, y) > 1 - r \text{ and } \nu_M(x, y) < r\}.$$

Definition 3.4. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. A subset A of X is said to be an open set (OS, for short) if for any $x \in A$ there exists $r \in]0, 1[$ such that $\mathcal{B}(x, r) \subseteq A$. The complement of an open set is called a closed set (CS, for short) in X .

Definition 3.5. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. Then

- (i) a sequence (x_n) in X is said to be convergent to a point $x \in X$ (i.e., $\lim_{n \rightarrow \infty} x_n = x$) if,

$$\lim_{n \rightarrow \infty} \mu_M(x_n, x) = 1, \lim_{n \rightarrow \infty} \sigma_M(x_n, x) = 1 \text{ and } \lim_{n \rightarrow \infty} \nu_M(x_n, x) = 0;$$

- (ii) a sequence (x_n) in X is said to be Cauchy sequence if, for each $k > 0$,

$$\lim_{n \rightarrow \infty} \mu_M(x_{n+k}, x_n) = 1, \lim_{n \rightarrow \infty} \sigma_M(x_{n+k}, x_n) = 1 \text{ and } \lim_{n \rightarrow \infty} \nu_M(x_{n+k}, x_n) = 0.$$

Definition 3.6. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. Then

- (i) if every Cauchy sequence is convergent, then X is said to be complete.
(ii) X is said to be compact if every sequence contains a convergent subsequence.

3.2. Fundamental properties of standard single valued neutrosophic metric space. In this section, we investigate the fundamental properties of standard single valued neutrosophic metric space that will be used in the following section.

Proposition 3.1. Every open ball in a standard single valued neutrosophic metric space $(X, M, *, \triangleleft, \diamond)$ is an open set.

Proof. Let $\mathcal{B}(x, r)$ be an open ball with radius r and center x , where $r \in]0, 1[$ and $x \in X$. Suppose that $y \in \mathcal{B}(x, r)$, this implies that

$$\mu_M(x, y) > 1 - r, \sigma_M(x, y) > 1 - r \text{ and } \nu_M(x, y) < r.$$

Let $r_0 = \mu_M(x, y)$. Then, there exist $s \in]0, 1[$ such that $r_0 > 1 - s > 1 - r$. Now, for a given r_0 and s such that $r_0 > 1 - s$. Then, Remark 3.2 guarantees that there exist $r_1, r_2, r_3 \in]0, 1[$ such that

$$r_0 * r_1 \geq 1 - s, r_0 \triangleleft r_2 \geq 1 - s \text{ and } (1 - r_0) \diamond (1 - r_3) \leq s.$$

Next, if we put $r_4 = \max\{r_1, r_2, r_3\}$ and consider the open ball $\mathcal{B}(y, 1 - r_4)$, then from the above, we can show that $\mathcal{B}(y, 1 - r_4) \subset \mathcal{B}(x, r)$ as follows:

Let $z \in \mathcal{B}(y, 1 - r_4)$. Then, $\mu_M(y, z) > r_4$, $\sigma_M(y, z) > r_4$ and $\nu_M(y, z) < 1 - r_4$. Furthermore, we obtain

$$\mu_M(x, z) \geq \mu_M(x, y) * \mu_M(y, z) \geq r_0 * r_4 \geq r_0 * r_1 \geq 1 - s > 1 - r,$$

$$\sigma_M(x, z) \geq \sigma_M(x, y) \triangleleft \sigma_M(y, z) \geq r_0 \triangleleft r_4 \geq r_0 \triangleleft r_2 \geq 1 - s > 1 - r$$

$$\text{and } \nu_M(x, z) \leq \nu_M(x, y) \diamond \nu_M(y, z) \leq (1 - r_0) \diamond (1 - r_4) \leq (1 - r_0) \diamond (1 - r_3) \leq s < r.$$

It follows that $z \in \mathcal{B}(x, r)$, and hence $\mathcal{B}(y, 1 - r_4) \subset \mathcal{B}(x, r)$. According to Definition 3.4, then it holds that $\mathcal{B}(x, r)$ is an open set. \square

Proposition 3.2. *Let $\mathcal{B}(x, r_1)$ and $\mathcal{B}(x, r_2)$ be two open balls with same center x in a standard fuzzy metric space $(X, M, *, \triangleleft, \diamond)$. Then either $\mathcal{B}(x, r_1) \subseteq \mathcal{B}(x, r_2)$ or $\mathcal{B}(x, r_2) \subseteq \mathcal{B}(x, r_1)$ where $r_1, r_2 \in]0, 1[$.*

Proof. Let $x \in X$ and $r_1, r_2 \in]0, 1[$. If $r_1 = r_2$, then $\mathcal{B}(x, r_1) = \mathcal{B}(x, r_2)$, and hence the result is trivially holds. Next, we assume that $r_1 \neq r_2$. Then, we distinguish two cases: $r_1 < r_2$ and $r_1 > r_2$.

- (i) If $r_1 < r_2$ and suppose that $y \in \mathcal{B}(x, r_1)$, then $\mu_M(x, y) > 1 - r_1$, $\sigma_M(x, y) > 1 - r_1$ and $\nu_M(x, y) < r_1$, which implies that $\mu_M(x, y) > 1 - r_2$, $\sigma_M(x, y) > 1 - r_2$ and $\nu_M(x, y) < r_2$. Therefore, $y \in \mathcal{B}(x, r_2)$, and hence $\mathcal{B}(x, r_1) \subseteq \mathcal{B}(x, r_2)$.
- (ii) If $r_1 > r_2$, then by applying a similar reasoning, we get $\mathcal{B}(x, r_2) \subseteq \mathcal{B}(x, r_1)$.

□

In the following theorem, we provide the topology induced by the single valued neutrosophic distance.

Theorem 3.1. *Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. Then it holds that the set*

$$\tau_M = \{A \subseteq X \mid x \in A \text{ if and only if there exists } r \in]0, 1[\text{ such that } \mathcal{B}(x, r) \subseteq A\}$$

is a topology on X and it is called the topology induced by the single valued neutrosophic set M .

Proof. (i) It is clear that $\emptyset, X \in \tau_M$.

- (ii) Consider $A_1, A_2 \in \tau_M$. Then, on the one hand, $A_1 \cap A_2 \subseteq X$. On the other hand, for any $x \in A_1$, there exists $r_1 \in]0, 1[$, such that $\mathcal{B}(x, r_1) \subseteq A_1$, and, for any $x \in A_2$, there exists $r_2 \in]0, 1[$, such that $\mathcal{B}(x, r_2) \subseteq A_2$. Here, we need to show that for any $x \in A_1 \cap A_2$, there exists $r \in]0, 1[$, such that $\mathcal{B}(x, r) \subseteq A_1 \cap A_2$. Indeed, due to Proposition 3.2, we distinguish two cases:

(a) If $\mathcal{B}(x, r_1) \subseteq \mathcal{B}(x, r_2)$ and take $r = r_1$, then $\mathcal{B}(x, r) \subseteq A_1 \cap A_2$.

(b) If $\mathcal{B}(x, r_2) \subseteq \mathcal{B}(x, r_1)$ and take $r = r_2$, then $\mathcal{B}(x, r) \subseteq A_1 \cap A_2$.

Hence, $A_1 \cap A_2 \in \tau_M$.

For the sake of completeness, we mention here, if we repeat this process for a finite number n ($n \in \mathbb{N}^*$) of sets $A_1, A_2, \dots, A_n \in \tau_M$, we deduce that $\bigcap_{1 \leq i \leq n} A_i \in \tau_M$.

- (iii) Suppose that $(A_i)_{i \in I} \in \tau_M$ and put $H = \bigcup_{i \in I} A_i$. We will show that $H \in \tau_M$.

Let $x \in H$, then $x \in \bigcup_{i \in I} A_i$. It follows that there exists $i_0 \in I$ such that $x \in A_{i_0}$.

The fact that $A_{i_0} \in \tau_M$ then, there exists $r_0 \in]0, 1[$ such that $\mathcal{B}(x, r_0) \subseteq A_{i_0}$.

Hence, $\mathcal{B}(x, r_0) \subseteq A_{i_0} \subseteq \bigcup_{i \in I} A_i = H$ which implies that $H \in \tau_M$.

Finally, we deduce that τ_M is a topology on X .

□

Remark 3.4. *Let (X, d) be an ordinary metric space and $(X, M, *, \triangleleft, \diamond)$ be the corresponding standard single valued neutrosophic metric space on X . Then, the topology τ_d induced by the metric d and the topology τ_M induced by the single valued neutrosophic set M are coincide. This means that $\tau_d = \tau_M$.*

The following theorem provides a basic property for the standard single valued neutrosophic metric spaces.

Theorem 3.2. *Every standard single valued neutrosophic metric space $(X, M, *, \triangleleft, \diamond)$ is a Hausdorff space.*

Proof. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space, and choose two distinct points x and y of X . Assume that $r_1 = \mu_M(x, y)$, $r_2 = \sigma_M(x, y)$ and $r_3 = \nu_M(x, y)$ with $r = \max\{r_1, r_2, 1 - r_3\}$. If we take $r_0 \in]r, 1[$, then, from Remark 3.2, it follows that there exist r_4, r_5, r_6 such that

$$r_4 * r_4 \geq r_0, r_5 \triangleleft r_5 \geq r_0 \text{ and } (1 - r_6) \diamond (1 - r_6) \leq 1 - r_0.$$

Next, if we put $r_7 = \max\{r_4, r_5, r_6\}$ and consider the open balls $\mathcal{B}(x, 1 - r_7)$ and $\mathcal{B}(y, 1 - r_7)$ then, it is clear that $B(x, 1 - r_7) \cap B(y, 1 - r_7) = \emptyset$. Now, if we suppose that there exists $z \in B(x, 1 - r_7) \cap B(y, 1 - r_7)$, then

$$r_1 = \mu_M(x, y) \geq \mu_M(x, z) * \mu_M(z, y) \geq r_7 * r_7 \geq r_4 * r_4 \geq r_0 > r_1,$$

$$r_2 = \sigma_M(x, y) \geq \sigma_M(x, z) \triangleleft \sigma_M(z, y) \geq r_7 \triangleleft r_7 \geq r_5 \triangleleft r_5 \geq r_0 > r_2,$$

and $r_3 = \nu_M(x, y) \leq \nu_M(x, z) \diamond \nu_M(z, y) \leq (1 - r_7) \diamond (1 - r_7) \leq (1 - r_6) \diamond (1 - r_6) \leq (1 - r_0) < r_3$. which is a contradiction. Hence, $(X, M, *, \triangleleft, \diamond)$ is a Hausdorff space. \square

Remark 3.5. We mention here that a Hausdorff space, also known as a separate space; is a topological space in which any two distinct points always admit disjoint neighborhoods.

3.3. Continuity property in standard single valued neutrosophic metric space.

In this subsection, we will study some interesting properties of single valued neutrosophic continuous mappings in standard single valued neutrosophic metric space. First, we introduce the notion of single valued neutrosophic continuous mapping.

Definition 3.7. Let $(X, M, *, \triangleleft, \diamond)$ and $(Y, M', *, \triangleleft, \diamond)$ be two standard single valued neutrosophic metric spaces. The mapping $f : X \rightarrow Y$ is a single valued neutrosophic continuous if and only if the inverse of each open set on Y is an open set on X .

Example 3.2. Let $(\mathbb{R}_+, M, *, \triangleleft, \diamond)$ and $(\mathbb{R}, M', *, \triangleleft, \diamond)$ be two standard single valued neutrosophic metric spaces. Define the t -norms $x * y = \min\{x, y\}$, $x \triangleleft y = \min\{x, y\}$ and the t -conorm $x \diamond y = \max\{x, y\}$, for all $x, y \in [0, 1]$. Define the single valued neutrosophic sets M and M' on \mathbb{R}_+^2 and \mathbb{R}^2 respectively as:

$$\begin{aligned} \mu_M(x, x') &= \frac{1}{e^{|x-x'|}} \quad \sigma_M(x, x') = \frac{1}{e^{|x-x'|}}, \text{ and } \nu_M(x, x') = \frac{|x-x'|}{1+|x-x'|}. \\ \mu_{M'}(y, y') &= \frac{1}{e^{|y-y'|}} \quad \sigma_{M'}(y, y') = \frac{1}{e^{|y-y'|}}, \text{ and } \nu_{M'}(y, y') = \frac{|y-y'|}{1+|y-y'|}. \end{aligned}$$

Then the mapping $f : (\mathbb{R}_+, M, *, \triangleleft, \diamond) \rightarrow (\mathbb{R}, M', *, \triangleleft, \diamond)$ defined by $f(x) = x + \alpha$, where $\alpha \in \mathbb{R}$ is a single valued neutrosophic continuous mapping. Indeed, let G be an open set on $(\mathbb{R}, M', *, \triangleleft, \diamond)$, and let $x \in f^{-1}(G)$, then it follows that $f(x) \in G$. Since G is an open set, then $\exists \varepsilon_{f(x)} \in]0, 1[$, where $\mathcal{B}_{\mathbb{R}}(f(x), \varepsilon_{f(x)}) \subseteq G$, and $f^{-1}(\mathcal{B}_{\mathbb{R}}(f(x), \varepsilon_{f(x)})) \subseteq f^{-1}(G)$. For $\varepsilon_x = \varepsilon_{f(x)}$, we will show that $f^{-1}(\mathcal{B}_{\mathbb{R}}(f(x), \varepsilon_{f(x)})) = \mathcal{B}_{\mathbb{R}_+}(x, \varepsilon_x)$. Let $x \in f^{-1}(\mathcal{B}_{\mathbb{R}}(f(x), \varepsilon_{f(x)}))$, it follows that $f(x) \in \mathcal{B}_{\mathbb{R}}(f(x), \varepsilon_{f(x)})$, and

$$\begin{aligned} \mu_{M'}(f(x), f(x')) &= \frac{1}{e^{|f(x)-f(x')|}} > 1 - \varepsilon_{f(x)}, \quad \sigma_{M'}(f(x), f(x')) = \frac{1}{e^{|f(x)-f(x')|}} > 1 - \varepsilon_{f(x)} \\ \text{and } \nu_{M'}(f(x), f(x')) &= \frac{|f(x)-f(x')|}{1+|f(x)-f(x')|} < \varepsilon_{f(x)}. \end{aligned}$$

Then, it holds that

$$\mu_{M'}(f(x), f(x')) = \frac{1}{e^{|(x+\alpha)-(x'+\alpha)|}} > 1 - \varepsilon_{f(x)}, \quad \sigma_{M'}(f(x), f(x')) = \frac{1}{e^{|(x+\alpha)-(x'+\alpha)|}} > 1 - \varepsilon_{f(x)}$$

$$\text{and } \nu_{M'}(f(x), f(x')) = \frac{|(x + \alpha) - (x' + \alpha)|}{1 + |(x + \alpha) - (x' + \alpha)|} < \varepsilon_{f(x)}.$$

This implies that

$$\mu_{M'}(f(x), f(x')) = \frac{1}{e^{|x-x'|}} > 1 - \varepsilon_{f(x)}, \quad \sigma_{M'}(f(x), f(x')) = \frac{1}{e^{|x-x'|}} > 1 - \varepsilon_{f(x)}$$

$$\text{and } \nu_{M'}(f(x), f(x')) = \frac{|x - x'|}{1 + |x - x'|} < \varepsilon_{f(x)}.$$

Hence,

$$\mu_M(x, x') > 1 - \varepsilon_x, \quad \sigma_M(x, x') > 1 - \varepsilon_x \text{ and } \nu_M(x, x') < \varepsilon_x.$$

We get that, $x \in \mathcal{B}_{\mathbb{R}_+}(x, \varepsilon_x)$. In the same way, we obtain the inverse implication. Therefore, $f^{-1}(\mathcal{B}_{\mathbb{R}}(f(x), \varepsilon_{f(x)})) = \mathcal{B}_{\mathbb{R}_+}(x, \varepsilon_x)$. We conclude that $f^{-1}(G)$ is an open set on $(\mathbb{R}_+, M, *, \triangleleft, \diamond)$, which satisfy the condition of the continuity property.

Remark 3.6. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space. Then the single valued neutrosophic identity mapping $Id : X \rightarrow X$ is a single valued neutrosophic continuous mapping.

Remark 3.7. Let $(X, M, *, \triangleleft, \diamond)$ and $(Y, M, *, \triangleleft, \diamond)$ be two standard single valued neutrosophic metric spaces. The mapping $f : X \rightarrow Y$ is a single valued neutrosophic continuous if and only if the inverse of each closed set on Y is a closed set on X .

The following proposition shows that the continuity property is preserved through the composition of single valued neutrosophic mappings.

Proposition 3.3. Let $(X, M, *, \triangleleft, \diamond)$, $(Y, M, *, \triangleleft, \diamond)$ and $(Z, M, *, \triangleleft, \diamond)$ be three standard single valued neutrosophic metric spaces and $f : X \rightarrow Y$, $g : Y \rightarrow Z$ be two continuous mappings. Then the composition $g \circ f : X \rightarrow Z$ is a continuous mapping.

Proof. Let A be an open set of Z . Definition 3.7 guarantees that $g^{-1}(A)$ is an open set of Y and by applying the same definition, we also get that $f^{-1}(g^{-1}(A))$ is an open set of X . Since $(g \circ f)^{-1}(A) = f^{-1}(g^{-1}(A))$, it follows that $g \circ f$ is a continuous mapping. \square

4. EDELSTEIN FIXED POINT THEOREM IN STANDARD SINGLE VALUED NEUTROSOPHIC METRIC SPACES

The aim of the present section, is to give an extension of Edelstein fixed point theorem in a standard single valued neutrosophic metric space. We start with the key results.

Lemma 4.1. Let $(X, M, *, \triangleleft, \diamond)$ be a standard single valued neutrosophic metric space and let (x_n) and (y_n) be two sequences in X . If $\lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$, then the following statements hold:

(i)

$$\begin{aligned} \mu_M(x, y) &\leq \liminf_{n \rightarrow \infty} \mu_M(x_n, y_n), \quad \sigma_M(x, y) \leq \liminf_{n \rightarrow \infty} \sigma_M(x_n, y_n) \\ \text{and } \nu_M(x, y) &\geq \limsup_{n \rightarrow \infty} \nu_M(x_n, y_n). \end{aligned} \quad (1)$$

(ii)

$$\begin{aligned} \mu_M(x, y) &\geq \limsup_{n \rightarrow \infty} \mu_M(x_n, y_n), \quad \sigma_M(x, y) \geq \limsup_{n \rightarrow \infty} \sigma_M(x_n, y_n) \\ \text{and } \nu_M(x, y) &\leq \liminf_{n \rightarrow \infty} \nu_M(x_n, y_n). \end{aligned} \quad (2)$$

Proof. (i) Let (x_n) and (y_n) be two sequences in X . From Definition 3.1 (iv), it follows that

$$\begin{aligned}\mu_M(x_n, y_n) &\geq \mu_M(x_n, x) * \mu_M(x, y) * \mu_M(y, y_n); \\ \sigma_M(x_n, y_n) &\geq \sigma_M(x_n, x) \triangleleft \sigma_M(x, y) \triangleleft \sigma_M(y, y_n); \\ \nu_M(x_n, y_n) &\leq \nu_M(x_n, x) \diamond \nu_M(x, y) \diamond \nu_M(y, y_n).\end{aligned}$$

Since $\lim_{n \rightarrow \infty} x_n = x$ and $\lim_{n \rightarrow \infty} y_n = y$, then it holds from Definition 3.5 that $\lim_{n \rightarrow \infty} \mu_M(x_n, x) = 1$, $\lim_{n \rightarrow \infty} \sigma_M(x_n, x) = 1$ and $\lim_{n \rightarrow \infty} \nu_M(x_n, x) = 0$. This implies that,

$$\begin{aligned}\liminf_{n \rightarrow \infty} \mu_M(x_n, y_n) &\geq 1 * \mu_M(x, y) * 1; \\ \liminf_{n \rightarrow \infty} \sigma_M(x_n, y_n) &\geq 1 \triangleleft \sigma_M(x, y) * 1; \\ \limsup_{n \rightarrow \infty} \nu_M(x_n, y_n) &\leq 0 \diamond \nu_M(x, y) \diamond 0.\end{aligned}$$

Hence,

$$\begin{aligned}\liminf_{n \rightarrow \infty} \mu_M(x_n, y_n) &\geq \mu_M(x, y); \\ \liminf_{n \rightarrow \infty} \sigma_M(x_n, y_n) &\geq \sigma_M(x, y); \\ \limsup_{n \rightarrow \infty} \nu_M(x_n, y_n) &\leq \nu_M(x, y).\end{aligned}$$

(ii) Follows from Definition 3.5 and (i). □

In the following theorem, we will establish the Edelstein fixed point theorem for standard single valued neutrosophic metric spaces.

Theorem 4.1 (Edelstein fixed point theorem). *Let $(X, M, *, \triangleleft, \diamond)$ be a compact standard single valued neutrosophic metric space. Let $T : X \rightarrow X$ be a mapping satisfying*

$$\begin{aligned}\mu_M(Tx, Ty) &> \mu_M(x, y), \sigma_M(Tx, Ty) > \sigma_M(x, y) \\ \text{and } \nu_M(Tx, Ty) &< \nu_M(x, y),\end{aligned}\tag{3}$$

for all $x \neq y$. Then, T has a unique fixed point.

Proof. Let $x \in X$ and define an iterative sequence (x_n) by $x_{n+1} = Tx_n$, for any $n \in \mathbb{N}$. Let us suppose that $x_n \neq x_{n+1}$ for each n (because, if $x_n = x_{n+1}$ then $x_n = Tx_n$, which means that x_n is a fixed point of T ; then the proof is finished). Therefore, $x_n \neq x_k$ where $n \neq k$. For otherwise, i.e., $x_n = x_k$ with $n \neq k$ we obtain

$$\begin{aligned}\mu_M(x_n, x_{n+1}) &= \mu_M(x_k, x_{k+1}) > \mu_M(x_{k-1}, x_k) > \dots \geq \mu_M(x_n, x_{n+1}), \\ \sigma_M(x_n, x_{n+1}) &= \sigma_M(x_k, x_{k+1}) > \sigma_M(x_{k-1}, x_k) > \dots \geq \sigma_M(x_n, x_{n+1}) \\ \text{and } \nu_M(x_n, x_{n+1}) &= \nu_M(x_k, x_{k+1}) < \nu_M(x_{k-1}, x_k) < \dots \leq \nu_M(x_n, x_{n+1})\end{aligned}$$

where $k > n$, a contradiction. Since X is compact, then (x_n) has a convergent subsequence (x_{n_i}) . Let $\lim_i x_{n_i} = y$. Also, we suppose that $y, Ty \notin (x_{n_i})$ where $i \in \mathbb{N}$ (if not, choose a subsequence with such a property). Due to the above hypotheses we now have

$$\mu_M(Tx_{n_i}, Ty) > \mu_M(x_{n_i}, y), \sigma_M(Tx_{n_i}, Ty) > \sigma_M(x_{n_i}, y)$$

$$\text{and } \nu_M(Tx_{n_i}, Ty) < \nu_M(x_{n_i}, y),$$

for all $i \in \mathbb{N}$. Then, by (1) we obtain

$$\lim \mu_M(Tx_{n_i}, Ty) > \lim \mu_M(x_{n_i}, y) = \mu_M(y, y) = 1,$$

$$\begin{aligned}\lim \sigma_M(Tx_{n_i}, Ty) &> \lim \sigma_M(x_{n_i}, y) = \mu_M(y, y) = 1 \\ \text{and } \lim \nu_M(Tx_{n_i}, Ty) &< \lim \nu_M(x_{n_i}, y) = \nu_M(y, y) = 0.\end{aligned}$$

Hence, $\lim Tx_{n_i} = Ty$. Similarly, we get

$$\lim T^2x_{n_i} = T^2y. \quad (4)$$

Now, recall that $Tx_{n_i} = Ty$ for all $i \in \mathbb{N}$. Notice that

$$\begin{aligned}\mu_M(x_{n_i}, Tx_{n_i}) &\leq \mu_M(Tx_{n_i}, T^2x_{n_i}) \leq \dots \leq \mu_M(x_{n_i}, Tx_{n_i}) \\ &\leq \mu_M(Tx_{n_i}, T^2x_{n_i}) \leq \dots \leq \mu_M(Tx_{n_{i+1}}, Tx_{n_{i+1}}) \\ &\leq \mu_M(Tx_{n_{i+1}}, T^2x_{n_{i+1}}) \leq \dots \leq 1, \\ \sigma_M(x_{n_i}, Tx_{n_i}) &\leq \sigma_M(Tx_{n_i}, T^2x_{n_i}) \leq \dots \leq \sigma_M(x_{n_i}, Tx_{n_i}) \\ &\leq \sigma_M(Tx_{n_i}, T^2x_{n_i}) \leq \dots \leq \sigma_M(Tx_{n_{i+1}}, Tx_{n_{i+1}}) \\ &\leq \sigma_M(Tx_{n_{i+1}}, T^2x_{n_{i+1}}) \leq \dots \leq 1\end{aligned}$$

and

$$\begin{aligned}\nu_M(x_{n_i}, Tx_{n_i}) &\geq \nu_M(Tx_{n_i}, T^2x_{n_i}) \geq \dots \geq \nu_M(x_{n_i}, Tx_{n_i}) \\ &\geq \nu_M(Tx_{n_i}, T^2x_{n_i}) \geq \dots \geq \nu_M(Tx_{n_{i+1}}, Tx_{n_{i+1}}) \\ &\geq \nu_M(Tx_{n_{i+1}}, T^2x_{n_{i+1}}) \geq \dots \geq 0.\end{aligned}$$

Therefore, the sequences formed from $\mu_M(x_{n_i}, Tx_{n_i})$ and $\mu_M(Tx_{n_i}, T^2x_{n_i})$ are convergent to a common limit [22] (we get the same if we replace μ_M by σ_M or ν_M). Thus, by (2), (4) and (1) we obtain

$$\begin{aligned}\mu_M(y, Ty) &\geq \limsup \mu_M(x_{n_i}, Tx_{n_i}) = \limsup \mu_M(Tx_{n_i}, T^2x_{n_i}) \\ &\geq \liminf \mu_M(Tx_{n_i}, T^2x_{n_i}) \geq \mu_M(Ty, T^2y), \\ \sigma_M(y, Ty) &\geq \limsup \sigma_M(x_{n_i}, Tx_{n_i}) = \limsup \sigma_M(Tx_{n_i}, T^2x_{n_i}) \\ &\geq \liminf \sigma_M(Tx_{n_i}, T^2x_{n_i}) \geq \sigma_M(Ty, T^2y), \\ \nu_M(y, Ty) &\leq \liminf \nu_M(x_{n_i}, Tx_{n_i}) = \liminf \nu_M(Tx_{n_i}, T^2x_{n_i}) \\ &\leq \limsup \nu_M(Tx_{n_i}, T^2x_{n_i}) \leq \nu_M(Ty, T^2y).\end{aligned}$$

Now suppose $y \neq Ty$. Then by (3), we have

$\mu_M(y, Ty) < \mu_M(Ty, T^2y)$, $\sigma_M(y, Ty) < \sigma_M(Ty, T^2y)$ and $\nu_M(y, Ty) > \nu_M(Ty, T^2y)$, which is a contradiction. (recall that μ_M and σ_M are continuous and non-decreasing, and, ν_M is continuous and non-increasing). Hence, $y = Ty$, i.e., y is a fixed point of T . Uniqueness follows at once from (3). \square

5. CONCLUSION

In this work, we have generalized the notion of standard fuzzy metric space introduced by J.R. Kider and Z.A. Hussain to the setting of single valued neutrosophic sets. Also, the fundamental properties related to the standard single valued neutrosophic metric space have been studied. Moreover, as an application we have been given an extension of the Edelstein fixed point theorem on standard single valued neutrosophic metric space. Future work is anticipated in multiple directions. We think it makes sense to study some other properties such as the connexity and the convexity of standard single valued neutrosophic metric space. Moreover, we intend to extend this work to other fixed point theorems.

REFERENCES

- [1] Abdel-Basset M., Chang V and Gamal A., (2019), Evaluation of the green supply chain management practices: A novel neutrosophic approach, *Computers in Industry*, 108, pp. 210–220.
- [2] Arkhangel'skii A.V. and Pontryagin L.S., (1990), *General Topology I: Basic Concepts and Constructions Dimension Theory*, Springer, 17, pp. 214–219.
- [3] Broumi S., Bakali A., Talea M. and Smarandache F., (2017), A Matlab Toolbox for interval valued neutrosophic matrices for computer applications, *Uluslararası Yönetim Bilişim Sistemleri ve Bilgisayar Bilimleri Dergisi*, 1, pp. 1–21.
- [4] Broumi S., Son L.H., Bakali A., Talea M. and Smarandache F., (2017), Selvachandran, G. Computing Operational Matrices in Neutrosophic Environments: A Matlab Toolbox, *Neutrosophic Sets and Systems*, 18, pp. 58–66.
- [5] Cheng H. D. and Guo Y., (2008), A new neutrosophic approach to image thresholding, *New Mathematics and Natural Computation*, 4, pp. 291–308.
- [6] Edelstein M., (1962), On fixed and periodic points under contractive mappings, *J. London Math. Soc.*, 37, pp. 74–79.
- [7] Etzion T. and Vardy A., Error-correcting codes in projective space, (2011), *IEEE Transaction on Information Theory*, 57, pp. 1165–1173.
- [8] Frank A. U., Campari I. and Formentini U., (1992), Theories and methods of spatio-temporal reasoning in geographic space; *International Conference GIS From Space to Territory*, *Lecture Notes in Computer Science*, 639, pp. 21–23.
- [9] Fréchet M., (1906), *Sur quelques points du calcul fonctionnel (Thèse)*, *Rendiconti del Circolo Matematico di Palermo*, 22, pp. 1–1774.
- [10] Garg H. and Nancy, Multiple attribute decision making based on immediate probabilities aggregation operators for single-valued and interval neutrosophic sets, *Journal of Applied Mathematics and Computing*, <https://doi.org/10.1007/s12190-020-01332-9>.
- [11] George A. and Veeramani P., (1994), On some results in fuzzy metric spaces, *Fuzzy Sets and Systems*, 64, pp. 395–399.
- [12] Hoffman K. and Kunze R., (1971), *Linear Algebra*, Prentice Hall, INC., Englewood Cliffs, New Jersey.
- [13] Kim J., Lim P.K., Lee J.G. and Hur K., (2018), Single valued neutrosophic relations, *Annals of Fuzzy Mathematics and Informatics*, 16, pp. 201–221.
- [14] Kider J.R. and Hussain Z.A., (2014), Continuous and Uniform Continuous Mappings on a Standard Fuzzy Metric Spaces, *Eng. and Tech. Journal*, 32, pp. 1111–1119.
- [15] Kirişçi M. and N. Şimşek, 2020, Neutrosophic metric spaces, *Mathematical Sciences*. <https://doi.org/10.1007/s40096-020-00335-8>.
- [16] Kirişçi M., N. Şimşek and Akyiğit M., 2020, Fixed point results for a new metric space, *Mathematical Methods in the Applied Sciences*, pp. 1–7. <https://doi.org/10.1002/mma.6189>.
- [17] Latreche A, Barkat O., Milles S. and Ismail F., (2020), Single valued neutrosophic mappings defined by single valued neutrosophic relations with applications, *Neutrosophic Sets and Systems*, 3, pp. 202–220.
- [18] Mondal K. and Pramanik S., (2015), Neutrosophic decision making model of school choice, *Neutrosophic Sets and Systems*, 7, pp. 62–68.
- [19] Okelo N.B., (2014), Properties of complete metric spaces and their applications to computer security, information and communication technology, *Evolving Trends in Engineering and Technology*, 1, pp. 23–28.
- [20] Pramanik S. and Roy T.K., (2014), Neutrosophic game theoretic approach to Indo-Pak conflict over Jammu-Kashmir, *Neutrosophic Sets and Systems*, 2, pp. 82–101.
- [21] Pramanik S. and Chackrabarti S. N., (2013), A study on problems of construction workers in West Bengal based on neutrosophic cognitive maps, *International Journal of Innovative Research in Science, Engineering and Technology*, 2, pp. 6387–6394.
- [22] Ray B.K., (1977), Some results on fixed points in metric and Banach spaces, *Bull. Acad. Polon. Math.*, 25, pp. 1243–1247.
- [23] Smarandache F., (1998), In: *Neutrosophy. Neutrosophic Property, Sets, and Logic*, American Research Press.
- [24] Smarandache F., (2007), In: *A Unifying Field in Logics: Neutrosophic Logic. Neutrosophy, Neutrosophic Set, Neutrosophic Probability and Statistics*, InfoLearnQuest.
- [25] Smarandache F., (2013), n-valued refined neutrosophic logic and its applications to Physics, *Progress in Physics*, 8, pp. 143–146.

- [26] Şimşek N. and Kirişçi M., (2019), Fixed point theorems in neutrosophic metric spaces, *Sigma J Eng. Nat. Sci.*, 10, pp. 221–230.
- [27] Şahin M. and Kargin A., (2018), Neutrosophic Triplet v -Generalized Metric Space, *Axioms*, 7, 67, doi:10.3390/axioms7030067.
- [28] Şahin M., Kargin A. and Yücel M., (2020), Neutrosophic Triplet Partial g -Metric Spaces, *Neutrosophic Sets and Systems*, 33, pp. 116–134.
- [29] Şahin M., Kargin A. and Uz M.S., (2020), Neutrosophic Triplet Partial Bipolar Metric Spaces, *Neutrosophic Sets and Systems*, 33, pp. 297–313.
- [30] Taş F., Topal S. and Smarandache F., Clustering Neutrosophic Data Sets and Neutrosophic Valued Metric Spaces, *Symmetry*, 10, pp. 430–442.
- [31] Walschap G., (2004), *Metric Structures in Differential Geometry*, Springer-Verlag.
- [32] Wang H, Smarandache F., Zhang Y.Q. and Sunderraman R., (2010), Single valued neutrosophic sets, *Multispace Multistruct*, 4, PP. 410–413.
- [33] Yang H.L., Guo Z.L. and Liao X., (2016), On single valued neutrosophic relations, *Journal of Intelligent and Fuzzy Systems*, 30, pp. 1045–1056.
- [34] Ye S. and Ye J., (2014), Dice similarity measure between single valued neutrosophic multisets and its application in medical diagnosis, *Neutrosophic Sets and Systems*, 6, pp. 49–54.
- [35] Ye J., (2014), Vector similarity measures of simplified neutrosophic sets and their application in multicriteria decision making, *International Journal of Fuzzy Systems*, 16, pp. 204–215.
- [36] Ye S., Fu J. and Ye J., (2014), Medical diagnosis sing distancebased similarity measures of single valued neutrosophic multisets, *Neutrosophic Sets and Systems*, 7, pp. 47–52.



Omar Barkat holds a BSc in Mathematics (1987) and a postgraduate degree in Applied Mathematics (2012). He benefited a research stay of 15 months (October 2016–December 2017) at the research unit KERMIT of Ghent University (Belgium), where he is a member in the research group on Fuzzy Mathematics and applications. Also, he holds a Doctorate in Mathematics (2019) and an Habilitation in Mathematics (2021). He is currently an associate professor and responsible for the training team (mathematics and computer science) at university center of Barika (Algeria). His main research interests include fuzzy set theory and its extensions with applications.



Soheyb Milles holds an MSc degree in Mathematics (2007), a postgraduate degree in Mathematical Logic (2010), a Doctorate in Mathematics (2017) and an Habilitation in Mathematics (2020). He is currently an associate professor at university center of Barika (Algeria), and he is the assistant director in charge of post-graduation, scientific research and external relations. His main research interests include lattice theory, fuzzy set theory and neutrosophic set theory.



Abdelkrim Latreche received the baccalaureate degree from 20 Aot 1955 University in 2013, and th Master and PhD degrees from 20 Aot 1955 university in 2015 and 2019 respectively. He is currently an associate professor at 20 Aot 1955 university. His research interests include fuzzy mathematics, and categorie theory.