

## INTEGRAL INEQUALITIES VIA $\log m$ -CONVEX FUNCTIONS

M. A. NOOR<sup>1</sup>, K. I. NOOR<sup>1</sup>, F. SAFDAR<sup>1</sup>, §

**ABSTRACT.** In this paper, we introduce and investigate a new concept of  $\log m$ -convex functions. We establish some new Hermite-Hadamard type integral inequalities via  $\log m$ -convex functions. Our results represent refinement and improvement of the previously known results. Several special cases are also discussed. The concept and technique of this paper may stimulate further research in this field.

**Keywords:**  $m$ -convex functions;  $\log m$ -convex functions; Hermite-Hadamard type inequalities.

**AMS Subject Classification:** 26D15; 26D10; 90C23

### 1. INTRODUCTION

Numerous applications in business, industry, art and medicine has turned convex analysis as one of the most interesting and useful field of mathematics for last few decades. The concept of convexity has played a crucial and significant part in the development of several branches of mathematical and engineering sciences. Consequently several new classes of convex functions and convex sets have been introduced and investigated, which make this area of research very attractive and useful, see [1, 2, 5, 6, 13, 14, 17, 20, 22]. It is well known that a function is convex if and only if, it satisfies the inequality which is known as Hermite-Hadamard inequality. It is one of the most important inequality, see [9, 10]. Several Hermite-Hadamard type inequalities have been derived for various classes of convex function using different techniques, see [7, 8, 11, 12, 15, 16, 17, 19, 20, 22].

Toader [23] introduced a new class of convex functions which is known as  $m$ -convex functions. This class is an intermediate form between the usual convexity and star shaped property. In this paper, we introduce a new class of convex functions relative to a constant  $m \in (0, 1]$ , which is known as  $\log m$ -convex function. It is worth mentioning that this class of  $\log m$ -convex functions is distinctly different from the  $\log m$ -convex functions which were considered in [3]. These  $\log m$ -convex functions are nonconvex functions.  $\log$  convex functions are of interest in many areas of mathematics and science. They play an important role in mathematical statistics and the theory of special functions, see [4, 21].

---

<sup>1</sup> Department of Mathematics, Comsats University, Islamabad, Pakistan.

e-mail: noormaslam@gmail.com; ORCID: <https://orcid.org/0000-0001-6105-2435>.

e-mail: khalidanoor@hotmail.com; ORCID: <https://orcid.org/0000-0002-5000-3870>.

e-mail: farhat.900@yahoo.com; ORCID: <https://orcid.org/0000-0002-0211-4891>.

§ Manuscript received: July 27, 2017; accepted: December 10, 2017.

TWMS Journal of Applied and Engineering Mathematics, Vol.10, No.4 © Işık University, Department of Mathematics, 2020; all rights reserved.

We derive some new Hermite-Hadamard integral inequalities for these nonconvex function. Our results include a wide class of known and new inequalities as special cases.

## 2. PRELIMINARIES

Let  $I$  be an interval in real line  $\mathbb{R}$ . Let  $f : I = [ma, b] \rightarrow \mathbb{R}$  be a continuous function and  $\eta(\cdot, \cdot) : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$  be a continuous bifunction. First of all, we recall the following well known results and concepts.

**Definition 2.1.** [23] A set  $\mathcal{S}$  is said to be  $m$ -convex set, if there exists a fixed constant  $m \in (0, 1]$ , such that

$$((1-t)ma + tb) \in \mathcal{S}, \quad \forall a, b \in \mathcal{S}, t \in [0, 1].$$

**Definition 2.2.** [23] A function  $f : I = [0, b] \rightarrow \mathbb{R}$  is said to be  $m$ -convex, where  $m \in (0, 1]$ , if

$$f(tma + (1-t)f(b)) \leq tmf(a) + (1-t)f(b), \quad \forall a, b \in I, t \in [0, 1].$$

It is denoted by  $K_m(b)$ , the class of all  $m$ -convex function on  $[0, b]$ .

If  $t = 1$ , then  $f(ma) \leq m(f)$ , which is called sub-homogeneous function.

We now introduce a new definition of log  $m$ -convex functions and derive various integral inequalities for these log  $m$ -convex functions.

**Definition 2.3.** A function  $f : I \rightarrow [0, \infty)$  is said to be log  $m$ -convex or multiplicatively  $m$ -convex if  $\log(f)$  is convex, or equivalently if for all  $a, b \in I$ ,  $m \in (0, 1)$  and  $t \in [0, 1]$ , one has the inequality

$$f(tma + (1-t)f(b)) \leq [mf(a)]^t [f(b)]^{1-t}, \quad \forall a, b \in I, t \in [0, 1]. \quad (1)$$

If  $t = \frac{1}{2}$  in (1), then

$$f\left(\frac{ma+b}{2}\right) \leq \sqrt{mf(a)[f(b)]}, \quad \forall a, b \in I. \quad (2)$$

which is known as Jensen log  $m$ -convex function.

From definition 2.3, we have

$$f(tma + (1-t)b) \leq [mf(a)]^t [f(b)]^{1-t} \leq tmf(a) + (1-t)f(b).$$

and

$$\log f(tma + (1-t)b) \leq t \log mf(a) + (1-t) \log f(b) \quad \forall a, b \in I, t \in [0, 1].$$

This means that log  $m$ -convex functions are  $m$ -convex functions. However the converse is not true.

It is worth mention that Bai et al. [3] introduced log $m$ -convex function as:

**Definition 2.4.** A function  $f : I \rightarrow [0, \infty)$  is said to be  $m$ -logarithmically convex if the inequality

$$f(ta + m(1-t)b) \leq [f(a)]^t [f(b)]^{m(1-t)}, \quad \forall a, b \in I, t \in [0, 1], \quad m \in (0, 1].$$

From definition 2.3 and Definition 2.4, it is clear that log  $m$ -convex functions are distinctly different from the log  $m$ convex functions defined by Bai et al. [3].

We will use the following notations throughout this paper.

- (1) Arithmetic Mean:  $A(a, b) = \frac{a+b}{2}, \quad \forall a, b \in \mathbb{R}_+.$
- (2) Geometric Mean:  $G(a, b) = \sqrt{ab}, \quad \forall a, b \in \mathbb{R}_+.$
- (3) Logarithmic mean:  $L(a, b) = \frac{b-a}{\log b - \log a}, \quad \forall a, b \in \mathbb{R}_+, a \neq b. \quad .$
- (4) Quadratic Mean:  $K(a, b) = \sqrt{\frac{a^2+b^2}{2}}, \quad \forall a, b \in \mathbb{R}_+..$

### 3. MAIN RESULTS

In this section, we establish several new integral inequalities of Hermite-Hadamard type for  $\log$   $m$ -convex functions.

**Theorem 3.1.** *Let  $f : I = [ma, b] \rightarrow (0, \infty)$  be a  $\log$   $m$ -convex function. Then*

$$\frac{1}{\sqrt{m}} f\left(\frac{ma+b}{2}\right) \leq \exp \frac{1}{(b-ma)} \int_{ma}^b \log f(x) dx \leq \sqrt{mf(a)f(b)}.$$

*Proof.* Let  $f$  be  $\log$   $m$ -convex function on  $I$ . Then

$$f(tma + (1-t)f(b)) \leq [mf(a)]^{1-t} [f(b)]^t, \tag{3}$$

which can be written as

$$\log f(tma + (1-t)f(b)) \leq (1-t) \log[mf(a)] + t \log[f(b)]. \tag{4}$$

Integrating (4) with respect to  $t$  on  $[0,1]$ , we have

$$\begin{aligned} \int_0^1 \log f(tma + (1-t)f(b)) dt &\leq \int_0^1 (1-t) \log[mf(a)] dt + \int_0^1 t \log[f(b)] dt \\ &= \log \sqrt{[mf(a)][f(b)]}. \end{aligned}$$

Thus

$$\frac{1}{b-ma} \int_{ma}^b \log f(x) dx \leq \log \sqrt{[mf(a)][f(b)]}. \tag{5}$$

Consider

$$\begin{aligned} f\left(\frac{ma+b}{2}\right) &= \frac{f((1-t)ma + tb + tma + (1-t)b)}{2} \\ &\leq \sqrt{[mf((1-t)ma + tb)][(f(tma + (1-t)b))]} \end{aligned} \tag{6}$$

From this it follows that

$$\log f\left(\frac{ma+b}{2}\right) \leq \frac{1}{2} \{ \log[mf((1-t)ma + tb)] + \log[f((tma + (1-t)b)] \}. \tag{7}$$

Integrating (7) with respect to  $t$  on  $[0,1]$ , we have

$$\begin{aligned} \log f\left(\frac{ma+b}{2}\right) &\leq \frac{1}{2} \left[ \frac{1}{(b-ma)} \int_{ma}^b \log mf(x)dx + \frac{1}{(ma-b)} \int_b^{ma} \log f(x)dx \right] \\ &= \left[ \frac{1}{2(b-ma)} \int_{ma}^b \log m dx + \frac{1}{(b-ma)} \int_{ma}^b \log f(x)dx \right] \\ &= \left[ \frac{\log m}{2(b-ma)}(b-ma) + \frac{1}{(b-ma)} \int_{ma}^b \log f(x)dx \right] \\ &= \left[ \log \sqrt{m} + \frac{1}{(b-ma)} \int_{ma}^b \log f(x)dx \right]. \end{aligned}$$

Thus

$$\log \frac{f\left(\frac{ma+b}{2}\right)}{\sqrt{m}} \leq \frac{1}{(b-ma)} \int_{ma}^b \log f(x)dx. \quad (8)$$

Combining(5) and (8), we have

$$\log \frac{f\left(\frac{ma+b}{2}\right)}{\sqrt{m}} \leq \frac{1}{(b-ma)} \int_{ma}^b \log mf(x)dx \leq \log \sqrt{mf(a)[f(b)]},$$

which implies that

$$\frac{1}{\sqrt{m}} f\left(\frac{ma+b}{2}\right) \leq \exp \frac{1}{(b-ma)} \int_{ma}^b \log f(x)dx \leq \sqrt{mf(a)f(b)}.$$

This completes the proof.  $\square$

**Corollary 3.1.** [7] *If  $m = 1$ , then under the assumption of Theorem 3.1, we have*

$$f\left(\frac{a+b}{2}\right) \leq \exp \frac{1}{(b-a)} \int_a^b \log f(x)dx \leq \sqrt{[f(a)][f(b)]},$$

*which is called the Hermite-Hadamard inequality for log convex function.*

**Theorem 3.2.** *Let  $f, g : I = [ma, b] \rightarrow (0, \infty)$  be log  $m$ -convex functions. Then*

$$\begin{aligned} &\frac{1}{b-ma} \int_{am}^b f(x)g(ma+b-x)dx \\ &\leq \frac{mg(a)f(b) + mf(a)g(b)}{2} \\ &\leq \frac{1}{2} \left\{ A[f(b), mf(a)]L[f(b), mf(a)] + A[mg(a), g(b)]L[mg(a), g(b)] \right\} \\ &\leq \frac{[mf(a) + (f(b))]^2}{16} + \frac{[mg(a) + g(b)]^2}{16} + \frac{mg(a)f(b) + mf(a)g(b)}{4}. \end{aligned}$$

*where  $A$  and  $L$  are Arithmetic and Logarithmic means respectively.*

*Proof.* Let  $f, g$  be log  $m$ -convex function. Then

$$\begin{aligned} f(tma + (1-t)b) &\leq [mf(a)]^t [f(b)]^{1-t} \\ g((1-t)ma + tb) &\leq [mg(a)]^{1-t} [g(b)]^t. \end{aligned}$$

Consider

$$\begin{aligned}
& \frac{1}{b-ma} \int_{ma}^b f(x)g(ma+b-x)dx \\
&= \int_0^1 f(tma+(1-t)b)g((1-t)ma+tb)dt \\
&\leq \int_0^1 \left[ \{mf(a)\}^t \{f(b)\}^{1-t} [\{mg(a)\}^{1-t} \{g(b)\}^t] \right] dt \\
&= [mg(a)f(b)] \int_0^1 \left[ \frac{mf(a)g(b)}{mg(a)f(b)} \right]^t dt \\
&= \frac{[mf(a)g(b) - mg(a)f(b)]}{\log [mf(a)g(b)] - \log [mg(a)f(b)]} \\
&\leq \frac{mg(a)f(b) + mf(a)g(b)}{2} \\
&\leq \frac{1}{2} \int_0^1 [\{f(tma+(1-t)b)\}^2 + \{g((1-t)ma+tb)\}^2] dt \\
&\leq \frac{1}{2} \int_0^1 \left\{ [\{mf(a)\}^t \{f(b)\}^{1-t}]^2 + [\{mg(a)\}^{1-t} \{g(b)\}^t]^2 \right\} dt \\
&= \frac{1}{2} \left\{ [f(b)]^2 \int_0^1 \left[ \frac{mf(a)}{f(b)} \right]^{2t} dt + [mg(a)]^2 \int_0^1 \left[ \frac{g(b)}{mg(a)} \right]^{2t} dt \right\} \\
&= \frac{1}{4} \left\{ \left[ \frac{[mf(a)]^2 - [f(b)]^2}{\log [mf(a)] - \log [f(b)]} \right] + \left[ \frac{[g(b)]^2 - [mg(a)]^2}{\log [g(b)] - \log [mg(a)]} \right] \right\} \\
&\leq \frac{1}{2} \left\{ A[f(b), mf(a)]L[f(b), mf(a)] + A[mg(a), g(b)]L[mg(a), g(b)] \right\} \\
&\leq \frac{1}{4} \int_0^1 [f(tma+(1-t)b) + g((1-t)ma+tb)]^2 dt \\
&\leq \frac{1}{4} \int_0^1 \left\{ [\{mf(a)\}^t \{f(b)\}^{1-t}]^2 + [\{mg(a)\}^{1-t} \{g(b)\}^t]^2 \right. \\
&\quad \left. + 2[\{mf(a)\}^t \{f(b)\}^{1-t}][\{mg(a)\}^{1-t} \{g(b)\}^t] \right\} dt \\
&= \frac{1}{4} \left\{ [f(b)]^2 \int_0^1 \left[ \frac{mf(a)}{f(b)} \right]^{2t} dt + [mg(a)]^2 \int_0^1 \left[ \frac{g(b)}{mg(a)} \right]^{2t} dt \right. \\
&\quad \left. + 2[mg(a)f(b)] \int_0^1 \left[ \frac{mf(a)g(b)}{mg(a)f(b)} \right]^t dt \right\} \\
&= \frac{1}{8} \left\{ \left[ \frac{[mf(a)]^2 - [f(b)]^2}{\log [mf(a)] - \log [f(b)]} \right] + \left[ \frac{[g(b)]^2 - [mg(a)]^2}{\log [g(b)] - \log [mg(a)]} \right] \right\} \\
&\quad + \frac{1}{2} \left\{ \frac{mf(a)g(b) - mg(a)f(b)}{\log [mf(a)g(b)] - \log [mg(a)f(b)]} \right\} \\
&\leq \frac{[mf(a) + f(b)]^2}{16} + \frac{[mg(a) + g(b)]^2}{16} + \frac{mg(a)f(b) + mf(a)g(b)}{4}.
\end{aligned}$$

This completes the proof.  $\square$

**Corollary 3.2.** [19] *If  $m = 1$ , then, under the assumption of Theorem 3.2, we have*

$$\begin{aligned} \frac{1}{b-a} \int_a^b f(x)g(a+b-x)dx &\leq \frac{g(a)f(b) + f(a)g(b)}{2} \\ &\leq \frac{1}{2} \left\{ A[f(b), f(a)]L[f(b), f(a)] + A[g(a), g(b)]L[g(a), g(b)] \right\} \\ &\leq \frac{[f(a) + f(b)]^2}{16} + \frac{[g(b) + g(a)]^2}{16} + \frac{g(a)f(b) + f(a)g(b)}{4}. \end{aligned}$$

**Corollary 3.3.** *If  $f = g$  and  $m = 1$ , then, under the assumption of Theorem 3.2, we obtain the result given in [24].*

**Theorem 3.3.** *Let  $f, g : I = [ma, b] \rightarrow (0, \infty)$  be log  $m$ -convex functions. If  $\alpha + \beta = 1$ , Then*

$$\begin{aligned} \frac{1}{b-ma} \int_{ma}^b f(x)g(ma+b-x)dx &\leq \alpha^2 \left[ \frac{[mf(a)]^{\frac{1}{\alpha}} - [f(b)]^{\frac{1}{\alpha}}}{[mf(a) - f(b)]} \right] L[f(b), mf(a)] \\ &\quad + \beta^2 \left[ \frac{[g(b)]^{\frac{1}{\beta}} - [mg(a)]^{\frac{1}{\beta}}}{[g(b) - mg(a)]} \right] L[mg(a), g(b)], \end{aligned}$$

where  $L$  is the Logarithmic mean.

*Proof.* Let  $f$  and  $g$  be log  $m$ -convex function on  $I$ . Then

$$\begin{aligned} f(tma + (1-t)b) &\leq [mf(a)]^t [f(b)]^{1-t} \\ g((1-t)ma + tb) &\leq [mg(a)]^{1-t} [g(b)]^t. \end{aligned}$$

Using young's inequality, that is,  $ab \leq \alpha a^{\frac{1}{\alpha}} + \beta b^{\frac{1}{\beta}}$ ,  $\forall \alpha, \beta > 0, \alpha + \beta = 1$ , we consider

$$\begin{aligned} &\frac{1}{b-ma} \int_{ma}^b f(x)g(ma+b-x)dx \\ &= \int_0^1 f(tma + (1-t)b)g((1-t)ma + tb)dt \\ &\leq \int_0^1 \left\{ \alpha [f(tma + (1-t)b)]^{\frac{1}{\alpha}} + \beta [g((1-t)ma + tb)]^{\frac{1}{\beta}} \right\} dt \\ &\leq \int_0^1 \left\{ \alpha \{ [mf(a)]^t [f(b)]^{1-t} \}^{\frac{1}{\alpha}} + \beta \{ [mg(a)]^{1-t} [g(b)]^t \}^{\frac{1}{\beta}} \right\} dt \\ &= \alpha [f(b)]^{\frac{1}{\alpha}} \int_0^1 \left[ \frac{mf(a)}{f(b)} \right]^{\frac{t}{\alpha}} dt + \beta [mg(a)]^{\frac{1}{\beta}} \int_0^1 \left[ \frac{g(b)}{mg(a)} \right]^{\frac{t}{\beta}} dt \\ &= \alpha^2 [f(b)]^{\frac{1}{\alpha}} \left[ \frac{(\frac{mf(a)}{f(b)})^u}{\log \frac{mf(a)}{f(b)}} \right]_0^1 + \beta^2 [mg(a)]^{\frac{1}{\beta}} \left[ \frac{(\frac{g(b)}{mg(a)})^u}{\log \frac{g(b)}{mg(a)}} \right]_0^1 \\ &= \alpha^2 \left[ \frac{[mf(a)]^{\frac{1}{\alpha}} - [f(b)]^{\frac{1}{\alpha}}}{[\log[mf(a)] - \log f(b)]} \right] + \beta^2 \left[ \frac{[g(b)]^{\frac{1}{\beta}} - [mg(a)]^{\frac{1}{\beta}}}{[\log[g(b)] - \log[mg(a)]]} \right] \\ &= \alpha^2 \left[ \frac{[mf(a)]^{\frac{1}{\alpha}} - [f(b)]^{\frac{1}{\alpha}}}{[mf(a) - f(b)]} \right] L[f(b), mf(a)] \\ &\quad + \beta^2 \left[ \frac{[g(b)]^{\frac{1}{\beta}} - [mg(a)]^{\frac{1}{\beta}}}{[g(b) - mg(a)]} \right] L[mg(a), g(b)]. \end{aligned}$$

This completes the proof. □

**Corollary 3.4.** [19] *If  $\alpha = \frac{1}{2}$ ,  $\beta = \frac{1}{2}$ , and  $m = 1$ , then, under the assumption of Theorem 3.3, we have*

$$\begin{aligned} \frac{1}{b-a} \int_a^b f(x)g(a+b-x)dx &\leq \frac{1}{4} \left\{ \left[ \frac{[f(a)]^2 - [f(b)]^2}{[f(a) - f(b)]} \right] L[f(b), f(a)] \right. \\ &\quad \left. + \left[ \frac{[g(b)]^2 - [g(a)]^2}{[g(b) - g(a)]} \right] L[g(a), g(b)] \right\}. \end{aligned}$$

**Corollary 3.5.** *If  $\alpha = \frac{1}{4}$ ,  $\beta = \frac{3}{4}$  and  $m = 1$ , then, under the assumption of Theorem 3.3, we have*

$$\begin{aligned} &\frac{1}{b-a} \int_a^b f(x)g(a+b-x)dx \\ &\leq \frac{1}{16} \left[ \frac{f^4(a) - f^4(b)}{f(b) - f(a)} \right] L[f(b), f(a)] + \frac{9}{16} \left[ \frac{g^{\frac{4}{3}}(b) - g^{\frac{4}{3}}(a)}{g(b) - g(a)} \right] L[g(a), g(b)]. \end{aligned}$$

**Theorem 3.4.** *Let  $f, g : I = [ma, b] \rightarrow (0, \infty)$  be increasing and  $\log$   $m$ -convex functions. Then*

$$\begin{aligned} 8L[mg(a), g(b)] \left\{ \frac{2}{m+1} f\left(\frac{ma+b}{2}\right) \right\} &\leq \frac{1}{b-ma} \int_{ma}^b f^4(x)dx + K^2[mg(a), g(b)] \\ &\quad A[mg(a), g(b)]L[mg(a), g(b)] + 8. \end{aligned}$$

*Proof.* Let  $f$  and  $g$  be  $\log$   $m$ -convex functions. Then

$$\begin{aligned} f(tma + (1-t)b) &\leq [mf(a)]^t [f(b)]^{1-t} \\ g((1-t)ma + tb) &\leq [mg(a)]^{1-t} [g(b)]^t. \end{aligned}$$

Using the inequality,

$$8xy \leq x^4 + y^4 + 8, \quad \forall x, y \in \mathbb{R},$$

we have

$$\begin{aligned} &8f(tma + (1-t)b)[mg(a)]^{1-t}t[g(b)]^t \\ &\leq f^4(tma + (1-t)b) + [mg(a)]^{4(1-t)}[g(b)]^t + 8. \end{aligned}$$

Now integrating the above inequality with respect to  $t$  on  $[0, 1]$ , we have

$$\begin{aligned} &8 \int_0^1 f(tma + (1-t)b)[mg(a)]^t [g(b)]^{1-t} dt \\ &\leq \int_0^1 f^4(tma + (1-t)b) dt + \int_0^1 [mg(a)]^{4t} [g(b)]^{4-4t} dt + 8. \end{aligned}$$

As  $f$  and  $g$  are increasing functions, we have

$$\begin{aligned} &8 \int_0^1 f(tma + (1-t)b) dt \int_0^1 [mg(a)]^t [g(b)]^{1-t} dt \\ &\leq \int_0^1 f^4(tma + (1-t)b) dt + \int_0^1 [mg(a)]^{4t} [g(b)]^{4-4t} dt + 8. \end{aligned}$$

From the above inequality, it is easy to observe that

$$\frac{8L[mg(a), g(b)]}{b - ma} \int_{ma}^b f(x) dx \leq \frac{1}{b - ma} \int_{ma}^b f^4(x) dx + K^2[mg(a), g(b)] A[mg(a), g(b)] L[mg(a), g(b)] + 8. \quad (10)$$

Now using the L.H.S of Hermite Hadamard's inequality in (10), we have

$$8L[mg(a), g(b)] \left\{ \frac{2}{m+1} f\left(\frac{ma+b}{2}\right) \right\} \leq \frac{1}{b - ma} \int_{ma}^b f^4(x) dx + K^2[mg(a), g(b)] A[mg(a), g(b)] L[mg(a), g(b)] + 8,$$

where A, L and K are Arithmetic, Logarithmic and Quadratic means respectively.  $\square$

**Corollary 3.6.** *If  $m = 1$ , then, under the assumption of Theorem 3.4, we have*

$$\begin{aligned} & 8L[g(a), g(b)] f\left(\frac{a+b}{2}\right) \\ & \leq \frac{1}{b-a} \int_a^b f^4(x) dx + K^2[g(a), g(b)] A[g(a), g(b)] L[g(a), g(b)] + 8. \end{aligned}$$

**Corollary 3.7.** *If  $f = g$  and  $m = 1$ , then, under the assumption of Theorem 3.2, we obtain the result given in [24].*

**Theorem 3.5.** *Let  $f, g : I = [ma, b] \rightarrow (0, \infty)$  be increasing and log  $m$ -convex functions. Then*

$$\begin{aligned} & \left(\frac{2}{m+1}\right) \left[ f\left(\frac{ma+b}{2}\right) L[mg(a), g(b)] + g\left(\frac{ma+b}{2}\right) L[f(b), mf(a)] \right] \\ & \leq \frac{1}{b - ma} \int_{ma}^b f(x)g(x) dx + L \left[ f(b)mg(a), [mf(a)g(b)] \right]. \end{aligned}$$

*Proof.* Let  $f$  and  $g$  be log  $m$ -convex functions. Then,  $\forall a, b \in I, t \in [0, 1]$ , we have

$$\begin{aligned} f(tma + (1-t)b) & \leq [mf(a)]^t [f(b)]^{1-t} \\ g((1-t)ma + tb) & \leq [mg(a)]^{1-t} [g(b)]^t. \end{aligned}$$

Using the inequality,

$$(a-b)(c-d) \geq 0. \quad \forall a, b, c, d \in \mathbb{R}, a < b, c < d,$$

we have

$$\begin{aligned} f(tma + (1-t)b) [mg(a)]^{1-t} [g(b)]^t & + [g((1-t)ma + tb)] [f(b)]^{1-t} [mf(a)]^t \\ & \leq f(tma + (1-t)b) g((1-t)ma + tb) \\ & + [mg(a)]^{1-t} [g(b)]^t [f(b)]^{1-t} [mf(a)]^t. \end{aligned}$$

Now integrating the above inequality with respect to  $t$  on  $[0, 1]$ , we have

$$\begin{aligned} & \int_0^1 [f(tma + (1-t)b)] [mg(a)]^{1-t} [g(b)]^t dt + \int_0^1 [g((1-t)ma + tb)] [f(b)]^{1-t} [mf(a)]^t dt \\ & \leq \int_0^1 [f(tma + (1-t)b)] g((1-t)ma + tb) dt + \int_0^1 [mg(a)]^{1-t} [g(b)]^t [f(b)]^{1-t} [mf(a)]^t dt. \end{aligned}$$



As  $f$  and  $g$  are increasing functions, we have

$$\begin{aligned} & \int_0^1 [f(tma + (1-t)b)]dt \int_0^1 [[mg(a)]^{1-t}[g(b)]^t] dt \\ & + \int_0^1 [g((1-t)ma + tb)]dt \int_0^1 [[f(b)]^{1-t}[mf(a)]^t] dt \\ & \leq \int_0^1 [f(tma + (1-t)b)]dt \int_0^1 [g((1-t)ma + tb)]dt \\ & + \int_0^1 [f(b)mg(a)]^{1-t}[mf(a)g(b)]^t dt. \end{aligned}$$

Now after some simple integration , we have

$$\begin{aligned} & \left(\frac{2}{m+1}\right) \left[ f\left(\frac{ma+b}{2}\right)L[mg(a), g(b)] + g\left(\frac{ma+b}{2}\right)L[f(b), mf(a)] \right] \\ & \leq \frac{1}{b-ma} \int_{ma}^b f(x)g(x)dx + L \left[ f(b)mg(a), [mf(a)g(b)] \right]. \end{aligned}$$

This completes the proof. □

**Corollary 3.8.** [24] *If  $m = 1$ , then, under the assumption of Theorem 3.5, we have*

$$\begin{aligned} & f\left(\frac{a+b}{2}\right)L[g(a), g(b)] + g\left(\frac{a+b}{2}\right)L[f(b), f(a)] \\ & \leq \frac{1}{b-a} \int_a^b f(x)g(x)dx + L[f(b)g(a), f(a)g(b)]. \end{aligned}$$

**Theorem 3.6.** *Let  $f, g : I = [ma, b] \rightarrow (0, \infty)$  be an increasing and  $\log m$ -convex functions. Then*

$$\begin{aligned} & \frac{1}{b-ma} \int_{ma}^b f^2(x)dx + A[f(b), mf(a)]L[f(b), mf(a)] + A[mg(a), g(b)] \\ & L[mg(a), g(b)] \\ & \geq \frac{2}{m+1} f\left(\frac{ma+b}{2}\right)L[f(b), mf(a)] + L[mg(a)f(b), mf(a)g(b)] \\ & + \frac{2}{m+1} f\left(\frac{ma+b}{2}\right)L[mg(a), g(b)]. \end{aligned}$$

*Proof.* Let  $f$  and  $g$  are  $\log$  - convex functions. Then

$$\begin{aligned} f(tma + (1-t)b) & \leq [mf(a)]^t[f(b)]^{1-t} \\ g((1-t)ma + tb) & \leq [mg(a)]^{1-t}[g(b)]^t. \end{aligned}$$

Using the inequality,

$$x^2 + y^2 + z^2 \geq xy + yz + zx. \quad \forall x, y, z \in \mathbb{R}, \tag{11}$$

we have

$$\begin{aligned} f^2(tma + (1-t)b) & + [mf(a)]^{2t}[f(b)]^{2(1-t)} + [mg(a)]^{2(1-t)}[g(b)]^{2t} \\ & \geq [f(tma + (1-t)b)][[mf(a)]^t[f(b)]^{1-t}] \\ & + [[mf(a)]^t[f(b)]^{1-t}][mg(a)]^{1-t}[g(b)]^t \\ & + [[mg(a)]^{1-t}[g(b)]^t]f(tma + (1-t)b). \end{aligned}$$

Now integrating the above inequality with respect to  $t$  on  $[0, 1]$ , we have

$$\begin{aligned}
 & \int_0^1 f^2(tma + (1-t)b)dt + \int_0^1 [[mf(a)]^{2t}[f(b)]^{2(1-t)}]dt \\
 & + \int_0^1 [[mg(a)]^{2(1-t)}[g(b)]^{2t}]dt \\
 & \geq \int_0^1 [f(tma + (1-t)b)[mf(a)]^t[f(b)]^{1-t}]dt \\
 & + \int_0^1 [[mf(a)]^t[f(b)]^{1-t}[mg(a)]^{1-t}[g(b)]^t]dt \\
 & + \int_0^1 [[mg(a)]^{1-t}[g(b)]^t f(tma + (1-t)b)]dt.
 \end{aligned} \tag{12}$$

To solve the integral in (12), let

$$\begin{aligned}
 \mathbb{A} & = \int_0^1 f^2(tma + (1-t)b)dt + \int_0^1 [[mf(a)]^{2t}[f(b)]^{2(1-t)}]dt \\
 & + \int_0^1 [[mg(a)]^{2(1-t)}[g(b)]^{2t}]dt \\
 & = \frac{1}{b-ma} \int_{ma}^b f^2(x)dx + A[f(b), mf(a)]\mathbb{L}[f(b), mf(a)] + A[mg(a), g(b)] \\
 & \quad L[mg(a), g(b)],
 \end{aligned}$$

and

$$\begin{aligned}
 \mathbb{B} & = \int_0^1 [f(tma + (1-t)b)[mf(a)]^t[f(b)]^{1-t}]dt \\
 & + \int_0^1 [[mf(a)]^t[f(b)]^{1-t}[mg(a)]^{1-t}[g(b)]^t]dt \\
 & + \int_0^1 [mg(a)]^{1-t}[g(b)]^t f(tma + (1-t)b)dt \\
 & \geq \frac{2}{m+1} f\left(\frac{ma+b}{2}\right)L[f(b), mf(a)] + L[mg(a)f(b), mf(a)g(b)] \\
 & + \frac{2}{m+1} f\left(\frac{ma+b}{2}\right)L[mg(a), g(b)].
 \end{aligned}$$

Substituting the values of  $\mathbb{A}$  and  $\mathbb{B}$  in (12), we have

$$\begin{aligned}
 & \frac{1}{b-ma} \int_{ma}^b f^2(x)dx + A[f(b), mf(a)]\mathbb{L}[f(b), mf(a)] + A[mg(a), g(b)] \\
 & L[mg(a), g(b)] \\
 & \geq \frac{2}{m+1} f\left(\frac{ma+b}{2}\right)L[f(b), mf(a)] + L[mg(a)f(b), mf(a)g(b)] \\
 & + \frac{2}{m+1} f\left(\frac{ma+b}{2}\right)L[mg(a), g(b)].
 \end{aligned}$$

This completes the proof.  $\square$

**Corollary 3.9.** *If  $m = 1$ , then, under the assumption of Theorem 3.6, we have*

$$\begin{aligned} & \frac{1}{b-a} \int_a^b f^2(x)dx + A[f(b), f(a)]L[f(b), f(a)] + A[g(a), g(b)]L[g(a), g(b)] \\ & \geq f\left(\frac{a+b}{2}\right)L[f(b), f(a)] + L[g(a)f(b), f(a)g(b)] + f\left(\frac{a+b}{2}\right)L[g(a), g(b)]. \end{aligned}$$

CONCLUSION

In this paper, we have introduced a new class of convex function relative to a constant  $m \in (0, 1]$ , which is known as  $\log m$ -convex function. New integral inequalities are obtained via these nonconvex functions. Some special cases are also discussed which have been obtained from our results. The technique of this paper may motivate new research.

ACKNOWLEDGEMENTS

The authors would like to thanks Rector, COMSATS University Islamabad, Pakistan, for providing excellent research and academic environments.

REFERENCES

- [1] Anderson, G.D., Vamanamurthy, and M. Vuorinen, M., (2007) Generalized convexity and inequalities, *J. Math. Anal. Appl.*, 335(2), pp. 1294-1308.
- [2] Alomari, M., Darus, M., and Dragomir, S. S., (2009), New inequalities of Simpson’s type for  $s$ -convex functions with applications, *RGMIA Res. Rep. Coll.*, 12 (4).
- [3] Bai, R. F., Qi, F. and Xi, B. Y., (2013), Hermite-Hadamard type inequalities for the  $m$  and  $(\alpha, m)$  logarithmically convex functions, *Filomat*, 27(1), pp. 1-7.
- [4] Carlson, B. C., (1977), *Special Functions of Applied Mathematics*, Academic Press, New York.
- [5] Cristescu, G. and Lupsa, L., (2002), *Non-connected Convexities and Applications*, Kluwer Academic Publishers, Dordrecht, Holland, (2002).
- [6] Cristescu, G., (2005), Improved integral inequalities for products of convex functions, *J. Inequal. Pure and Appl. Math.*, 6(2), Article: 35.
- [7] Dragomir, S. S. and Mond, B., (1998), Integral inequalities of Hadamard type for  $\log$ -convex functions, *Demonstratio*, 31, pp. 354-364.
- [8] Dragomir, S. S. and Pearce, C. E. M., (2000), *Selected topics on Hermite-Hadamard inequalities and applications*, Victoria University, Australia, (2000).
- [9] Hadamard, J., (1893), Etude sur les proprietes des fonctions entieres e.t en particulier dune fonction consideree par Riemann, *J. Math. Pure. Appl.*, 58, pp. 171-215.
- [10] C. Hermite, C., (1883), Sur deux limites d’une integrale definie, *Mathesis*, 3, pp. 82.
- [11] Hyers, D. H., and Ulam, S. M., (1952), Approximately convex functions, *Proc. Amer. Math. Soc.*, 3, 821-828, (1952).
- [12] Miheșan, V. G., (1993), A generalization of the convexity, *Seminar on Functional Equations, Approximation and Convex*, Cluj-Napoca (Romania), (1993).
- [13] Niculescu, C. P. and Persson, L. E., (2018), *Convex Functions and Their Applications*. Springer-Verlag, New York.
- [14] Niculescu, C. P., (2012), The Hermite-Hadamard inequality for  $\log$ -convex functions, *Nonlinear Analysis*, 75, pp. 662-669.
- [15] Noor, M. A., Qi, F. and Awan, M. U., (2013), Hermite-Hadamard type inequalities for  $\log$ - $h$ -convex functions, *Analysis*, 33, pp. 367-375.
- [16] M. A., Noor, K. I. and Awan, M. U., (2014), Hermite-Hadamard inequalities for relative semi-convex functions and applications, *Filomat*, 28(2), pp. 221-230.
- [17] M. A., Noor, K. I. and Awan, M. U., (2015), Generalized convexity and integral inequalities, *Appl. Math. Inf. Sci.*, 9(1), pp. 233-243.
- [18] Noor, M. A., (2007), On Hadamard integral inequalities involving two  $\log$ -preconvex functions, *J. Inequal. Pure and App. Math.*, 8(3).
- [19] M. A., Noor, K. I. and Awan, M. U., (2014), Some characterizations of harmonically  $\log$ -convex functions, *Proc. Jangjeon. Math. Soc.*, 17, pp. 51-61.

- [20] M. A., Noor, K. I., Iftikhar, S. and F. Safdar, F.,(2015), Integral inequaities for relative harmonic  $(s,\eta)$ -convex functions, Appl. Math. Inf. Sci, 1(1), pp. 27-34.
- [21] Pecaric, J. E., Proschan, F. and Y.L. Tong, Y. L.,(1992), Convex functions, partial orderings and statistical applications, Academic Press, Boston.
- [22] M. A., Noor, K. I. and Safdar, F., (2017), Integral inequaities via  $(\alpha,m)$ -convex functions, J. Nonlinear. Func. Anal, 2017, pp.
- [23] G. Toader, G., (1985), Some generalizations of the convexity, Proceedings of the Colloquium on Approximation and Optimization, Univ. Cluj-Napoca, pp. 329-338.
- [24] M. Tunc, M.,(2014), Some integral inequaities for logarithmically convex functions, J. Egypt. Math. Soc, 22, pp. 177-181.



**Muhammad Aslam Noor** earned his PhD degree from Brunel University, UK. His field of interest and specializations are: Variational Inequalities, Optimization and Numerical Analysis. He has been awarded President's Award for pride of performance and Sitar-i-Imtiaz). He has more than 1030 research papers to his credit. Dr. Noor is a Highly Cited Researcher in Mathematics



**Khalida Inayat Noor** obtained her PhD from Wales University (UK). She was awarded HEC best research paper award and CIIT Medal for innovation in 2009. Professor Khalida Noor was awarded President's Award for pride of performance. Her field of interest and specializations are Geometric function theory and Convex analysis. She has published more than 680 research articles in reputed mathematical and engineering journals and is member of editorial boards of several international journals.



**Dr. Farhat Safdar** earned her PhD degree from COMSATS University Islamabad, Islamabad, Pakistan. Dr. Farhat Safdar is a faculty member of Department of Mathematics, SBK Women University, Quetta, Pakistan. She is the author of several research articles. Her field of interest is Integral Inequalities and Convex Analysis.

---

---