

## COLOR LAPLACIAN ENERGY OF SOME CLUSTER GRAPHS

S. D'SOUZA<sup>1</sup>, GOWTHAM H. J.<sup>1\*</sup>, PRADEEP G. BHAT<sup>1</sup>, GIRIJA K. P.<sup>1</sup>, §

ABSTRACT. The color energy of a graph  $G$  is defined as the sum of the absolute values of the color eigenvalues of  $G$ . The graphs with large number of edges are referred as cluster graphs. Cluster graphs are obtained from complete graphs by deleting few edges according to some criteria. Bipartite cluster graphs are obtained by deleting few edges from complete bipartite graphs according to some rule. In this paper, we study the color Laplacian energy of cluster graphs and bipartite cluster graphs obtained by deleting the edges of complete and complete bipartite graph respectively.

Keywords: color Laplacian matrix, color Laplacian eigenvalues, color Laplacian energy

AMS Subject Classification: 05C15, 05C50

### 1. INTRODUCTION

The energy  $E(G)$  of a graph  $G$ , defined as the sum of the absolute values of its eigenvalues, belongs to the most popular graph invariants in chemical graph theory. It originates from the  $\pi$ -electron energy in the Hückel molecular orbital model, but has also gained purely mathematical interest. Gutman introduced this definition of the energy of a simple graph in his paper “The energy of a graph” [9]. In the past decade, interest in graph energy has increased and many different versions have been introduced. In 2006, Gutman and Zhou defined the Laplacian energy of a graph as the sum of the absolute deviations (i.e., distance from the mean) of the eigenvalues of its Laplacian matrix [10]. Similar variants of graph energy were developed for the signless Laplacian matrix, the distance matrix, the incidence matrix, and even for a general matrix not associated with a graph [1].

Let  $G$  be a simple undirected graph with  $n$  vertices. Let  $A(G)$  be the adjacency matrix of  $G$  and  $D(G)$  be the diagonal matrix of vertex degrees. Then the Laplacian matrix of

---

<sup>1</sup> Department of Mathematics, Manipal Institute of Technology, Academy of Higher Education, Manipal, Karnataka, 576104, India.

e-mail: sabitha.dsouza@manipal.edu; ORCID: [http://orcid.org/0000 0002 2728 6403](http://orcid.org/0000_0002_2728_6403).

e-mail: gowtham.hj@manipal.edu; ORCID: [http://orcid.org/ 0000 0001 5276 2363](http://orcid.org/0000_0001_5276_2363).

\* Corresponding author.

e-mail: pg.bhat@manipal.edu; ORCID: [http://orcid.org/0000 0003 2179 6207](http://orcid.org/0000_0003_2179_6207).

e-mail: girijakp16@gmail.com; ORCID: [http://orcid.org/0000 0003 4236 602x](http://orcid.org/0000_0003_4236_602x).

§ Manuscript received: January 09, 2020; accepted: April 13, 2020.

TWMS Journal of Applied and Engineering Mathematics, Vol.12, No.1 © Işık University, Department of Mathematics, 2022; all rights reserved.

$G$  is  $L(G) = D(G) - A(G)$ . The Laplacian energy of graph  $G$  [10] is defined as  $LE(G) = \sum_{i=1}^n \left| \mu_i - \frac{2m}{n} \right|$ , where  $m$  is the number of edges in the graph  $G$ .

A coloring of a graph  $G$  [12] is a coloring of its vertices such that no two adjacent vertices share the same color. The minimum number of colors needed for the coloring of a graph  $G$  is called the chromatic number of  $G$  and is denoted by  $\chi(G)$ .

In 2013, C. Adiga, E. Sampathkumar, M. A. Sriraj and A. S. Shrikanth [2], have introduced the energy of colored graph. The entries of the color adjacency matrix  $A_c(G)$  are as follows: If  $c(v_i)$  is the color of vertex  $v_i$ , then

$$a_{ij} = \begin{cases} 1, & \text{if } v_i \text{ and } v_j \text{ are adjacent with } c(v_i) \neq c(v_j), \\ -1, & \text{if } v_i \text{ and } v_j \text{ are non-adjacent with } c(v_i) = c(v_j), \\ 0, & \text{otherwise.} \end{cases}$$

The color energy of a graph  $G$  is the sum of absolute values of the color eigenvalues of  $G$ , i. e.,  $E_c(G) = \sum_{i=1}^n |\lambda_i|$ , where  $\lambda_1, \lambda_2, \dots, \lambda_n$  are the color eigenvalues of graph  $G$ .

In 2015, P. G. Bhat and S. D'Souza [7] have introduced the color Laplacian matrix of  $G$  as  $L_c(G) = D(G) - A_c(G)$ . The eigenvalues  $\mu_1, \mu_2, \dots, \mu_n$  are called color Laplacian eigenvalues of the graph  $G$ . The characteristic polynomial of color Laplacian matrix of  $G$  is denoted by  $\phi(L_c(G), \mu)$ . The color Laplacian energy of graph is defined as  $LE_c(G) = \sum_{i=1}^n \left| \mu_i - \frac{2m}{n} \right|$ , where  $m$  is the number of edges in the graph  $G$ . For more information on energy, Laplacian energy and color energy of a graph we refer [1, 3, 4, 5, 6, 8].

I. Gutman and L. Pavlović [11] introduced cluster graphs obtained by deleting the edges of complete graphs and found energies of cluster graphs. H. B. Walikar and H. S. Ramane [14] introduced bipartite cluster graphs obtained by deleting edges of complete bipartite graph. In section 2, we establish color Laplacian energy of some cluster graphs. In section 3, we establish color Laplacian energy of bipartite cluster graphs.

## 2. COLOR LAPLACIAN ENERGY OF SOME CLUSTER GRAPHS

**Definition 2.1.** [13] Let  $(K_m)_i, i = 1, 2, \dots, k, 1 \leq k \leq \lfloor \frac{n}{m} \rfloor, 1 \leq m \leq n$ , be independent complete subgraphs with  $m$  vertices of the complete graph  $K_n, n \geq 3$ . The cluster graph  $Ka_n(m, k)$  is obtained from  $K_n$ , by deleting all edges of  $(K_m)_i, i = 1, 2, \dots, k$ . In addition  $Ka_n(m, 0) = Ka_n(0, k) = Ka_n(0, 0) = K_n$ .

**Example 2.1.** Consider the cluster graph




FIGURE 1. Graph  $Ka_5(2, 2)$

Let  $c(v_1) = c(v_2) = 1, c(v_3) = c(v_4) = 2, c(v_5) = 3$ . The color Laplacian matrix is,

$$L_c(Ka_5(2, 2)) = \left[ \begin{array}{cccc|c} 3 & 1 & -1 & -1 & -1 \\ 1 & 3 & -1 & -1 & -1 \\ -1 & -1 & 3 & 1 & -1 \\ -1 & -1 & 1 & 3 & -1 \\ -1 & -1 & -1 & -1 & 4 \end{array} \right] = \left[ \begin{array}{c|c} X_{4 \times 4} & -J_{4 \times 1} \\ \hline -J_{1 \times 4} & 5I_1 - J_{1 \times 1} \end{array} \right].$$

Where  $X = \left[ \begin{array}{c|c} 2I_2 + J_2 & -J_2 \\ \hline -J_2 & 2I_2 - J_2 \end{array} \right]$ ,  $I$  is identity matrix and  $J$  is the matrix with all entries one.

**Theorem 2.1.** For  $n \geq 3$ ,  $0 \leq k \leq \lfloor \frac{n}{m} \rfloor$  and  $1 \leq m \leq n$ ,

$$LE_c(Ka_n(m, k)) = \frac{1}{n} \left[ 2km(m-1)(n-1) - 2mk^2(m-1)^2 + n(n-m-1 + \sqrt{m^2(1-4k) + (n-1)^2 + 2m(n-1+2k)}) \right].$$

*Proof.* Consider the cluster graph  $Ka_n(m, k)$  obtained from complete graph  $K_n$  by deleting all edges of  $(K_m)_i$ ,  $i = 1, 2, \dots, k$ . Since  $\chi(Ka_n(m, k)) = n - (m-1)k$ , we have

$$L_c(Ka_n(m, k)) = \left[ \begin{array}{c|c} X_{mk \times mk} & -J_{mk \times (n-mk)} \\ \hline -J_{(n-mk) \times mk} & (nI - J)_{(n-mk) \times (n-mk)} \end{array} \right]$$

$$X = \left[ \begin{array}{c|c|c|c} (n-m-1)I_m + J_m & -J_m & \dots & -J_m \\ \hline -J_m & (n-m-1)I_m + J_m & \dots & -J_m \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline -J_m & -J_m & \dots & (n-m-1)I_m + J_m \end{array} \right]_{mk \times mk}$$

Consider  $\det(\mu I - L_c(Ka_n(m, k)))$ .

Step 1: Replace  $R_i$  by  $R'_i = R_i - R_{i+1}$ , for  $i = 1, 2, \dots, m-1, m+1, m+2, \dots, 2m-1, 2m+1, \dots, mk-2, mk-1, mk+1, mk+2, \dots, n-2, n-1$ . Then,  $\det(\mu I - L_c(Ka_n(m, k)))$  will reduce to a new determinant, say  $\det(C)$ .

Step 2: In  $\det(C)$ , replacing  $C_i$  by  $C'_i = C_i + C_{i-1}$ , for  $i = 2, 3, \dots, m, m+2, m+3, \dots, 2m, 2m+2, \dots, mk-1, mk, mk+2, mk+3, \dots, n-1, n$ , a new determinant  $\det(D)$  is obtained.

Step 3: On expanding  $\det(D)$  along the rows  $R_i$ , for  $i = 1, 2, 3, \dots, k-1, k+1, k+2, \dots, 2k-1, 2k+1, 2k+2, \dots, mk-2, mk-1, mk+1, mk+2, \dots, n-2, n-1$ , it becomes  $(\mu - n + m + 1)^{(m-1)k} (\mu - n)^{n-mk-1} \det(E)$ . Where

$$\det(E) = \begin{vmatrix} \mu - n + 1 & m & m & \dots & m & n - mk \\ m & \mu - n + 1 & m & \dots & m & n - mk \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ m & m & m & \dots & \mu - n + 1 & n - mk \\ m & m & m & \dots & m & \mu - mk \end{vmatrix}_{(k+1) \times (k+1)}$$

Step 4: In  $\det(E)$ , replacing  $R_i$  by  $R'_i = R_i - R_{i+1}$ , for  $i = 1, 2, \dots, k$ , it reduces to

$$\det(E) = \begin{vmatrix} \mu - n - m + 1 & -(\mu - n - m + 1) & 0 & \dots & 0 & 0 \\ 0 & \mu - n - m + 1 & -(\mu - n - m + 1) & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & 0 & \dots & \mu - n - m + 1 & n - \mu \\ 0 & 0 & 0 & \dots & m & \mu - mk \end{vmatrix}$$

Step 5: In  $\det(E)$ , replacing  $C_i$  by  $C'_i = C_i + C_{i-1} + \dots + C_1$ , for  $i = k, k - 1, \dots, 2$  it simplifies to

$$\begin{aligned} \det(E) &= (\mu - n - m + 1)^{k-1} \begin{vmatrix} \mu - n - m + 1 & n - \mu \\ mk & \mu + mk \end{vmatrix} \\ &= (\mu - n - m + 1)^{k-1} (\mu^2 - (m + n - 1)\mu - (m^2 - m)k). \end{aligned}$$

Thus,

$$\begin{aligned} \phi(L_c(Ka_n(m, k), \mu)) &= (\mu - n + m + 1)^{(m-1)k} (\mu - n)^{n-mk-1} (\mu - n - m + 1)^{k-1} \\ &\quad (\mu^2 - (m + n - 1)\mu - (m^2 - m)k). \end{aligned}$$

So, the color Laplacian spectrum of  $Ka_n(m, k)$  is  $\{n + m - 1$  ( $k - 1$  times),  $n - m - 1$  ( $(m - 1)k$  times),  $n$  ( $n - mk - 1$  times),

$$\left. \frac{m + n - 1 + \sqrt{(m + n - 1)^2 - 4(m^2 - m)k}}{2}, \frac{m + n - 1 - \sqrt{(m + n - 1)^2 - 4(m^2 - m)k}}{2} \right\}.$$

Average degree of  $Ka_n(m, k)$  is  $\frac{n(n - 1) - km(m - 1)}{n}$ .

Hence,

$$\begin{aligned} LE_c(Ka_n(m, k)) &= (m - 1)k \left| (n - m - 1) - \frac{n(n - 1) - km(m - 1)}{n} \right| + (n - mk - 1) \\ &\quad \left| n - \frac{n(n - 1) - km(m - 1)}{n} \right| + (k - 1) |(n + m - 1) - \\ &\quad \frac{n(n - 1) - km(m - 1)}{n}| + \\ &\quad \left| \frac{(m + n - 1) + \sqrt{(m + n - 1)^2 - 4k(m^2 - m)}}{2} - \right. \\ &\quad \left. \frac{n(n - 1) - km(m - 1)}{n} \right| \\ &\quad + \left| \frac{(m + n - 1) - \sqrt{(m + n - 1)^2 - 4k(m^2 - m)}}{2} - \right. \\ &\quad \left. \frac{n(n - 1) - km(m - 1)}{n} \right| \\ &= \frac{1}{n} \left[ 2km(m - 1)(n - 1) - 2mk^2(m - 1)^2 + n(n - m - 1) + \right. \\ &\quad \left. \sqrt{m^2(1 - 4k) + (n - 1)^2 + 2m(n - 1 + 2k)} \right]. \end{aligned}$$

□

**Definition 2.2.** [11] For fixed integers  $n \geq 3$  and  $0 \leq k \leq \lfloor \frac{n}{2} \rfloor$ , the cluster graph  $K_{b_n}(k)$  is obtained from  $K_n$  by the deletion of  $k$  independent edges.

**Corollary 2.1.** For  $n \geq 3$  and  $0 \leq k \leq \lfloor \frac{n}{2} \rfloor$ ,

$$LE_c(K_{b_n}(k)) = \frac{1}{n} \left[ 4k(n - 4k - 1) + n \left( n - 3 + \sqrt{(n - 1)^2 + 4(n - 2k)} \right) \right].$$

*Proof.* Observe that  $K_{b_n}(k)$  is a special case of  $Ka_n(m, k)$ , when  $m=2$ . Thus, by substituting  $m = 2$  in Theorem 2.1, the result follows. □

**Definition 2.3.** [11] Let  $n \geq 3$  and  $k$ , where  $1 \leq k \leq n - 1$ , be fixed integers. The cluster graph  $K_{c_n}(k)$  is obtained from  $K_n$  by deleting a  $k$ -clique.

**Corollary 2.2.** For  $n \geq 3$  and  $1 \leq k \leq n - 1$ ,

$$LE_c(K_{c_n}(k)) = \frac{1}{n} \left[ 2k(k - 1)(n - k) + n(n - k - 1 + \sqrt{(n - 1)^2 - 3k^2 + 2k(n + 1)}) \right].$$

*Proof.* The proof follows by noting that  $K_{c_n}(k) = Ka_n(m, 1)$  in Theorem 2.1. □

**Definition 2.4.** [11] Let  $n \geq 3$  and  $1 \leq k \leq \lfloor \frac{n}{3} \rfloor$  be fixed integers. The cluster graph  $K_{f_n}(k)$  is obtained from  $K_n$  by the deletion of  $k$  disjoint triangles.

**Corollary 2.3.** For  $n \geq 3$  and  $1 \leq k \leq \lfloor \frac{n}{3} \rfloor$ ,

$$LE_c(K_{f_n}(k)) = \frac{1}{n} \left[ 12k(n - 2k - 1) + n \left( n - 4 + \sqrt{(n - 1)^2 + 3(2n - 8k + 1)} \right) \right].$$

*Proof.* The proof follows by noting that  $K_{f_n}(k) = Ka_n(3, k)$  in Theorem 2.1. □

**Definition 2.5.** [11] For fixed integers  $n$  and  $k$ ,  $n \geq 3$  and  $0 \leq k \leq n - 1$ , the graph  $K_{a_n}(k)$  is obtained from  $K_n$  by the deletion of  $k$  edges with a common end vertex.

**Theorem 2.2.** For  $n \geq 3$  and  $0 \leq k \leq n - 1$ ,

$$\phi(L_c(K_{a_n}(k), \mu)) = (\mu - n)^{n-k-2}(\mu - n + 1)^{k-2}(\mu^4 - (3n - k - 2)\mu^3 + ((3n^2 - 4n) - (2n - 1)k)\mu^2 - ((n^3 - 2n^2 + 2n - 3) - (n^2 - n + 2)k)\mu + (2n^2 - 6n + 3) - (2n - 3)k).$$

*Proof.* Let  $v_1, v_2, \dots, v_n$  be the vertices of complete graph  $K_n$ . The graph  $K_{a_n}(k)$  is obtained from  $K_n$  by the deletion of  $k$  edges with common end vertex  $v_i$ , where  $i = 1, 2, 3, \dots, n$ . and  $\chi(K_{a_n}(k)) = n - 1$ . Then, we have

$$L_c(K_{a_n}(k)) = \left[ \begin{array}{c|c|c} (n - k - 1)_1 & C_{1 \times k} & -J_{1 \times n - k - 1} \\ \hline C_{k \times 1}^T & (n - 1)I_k - J_k & -J_{k \times n - k - 1} \\ \hline -J_{n - k - 1 \times 1} & -J_{n - k - 1 \times k} & (nI - J)_{n - k - 1} \end{array} \right]$$

Where  $C = [1 \ 0 \ 0 \dots 0]_{1 \times k}$ . Consider  $\det(\mu I - L_c(K_{a_n}(k)))$ .

Step 1: Replace  $R_i$  by  $R'_i = R_i - R_{i+1}$ , for  $i = 2, 3, \dots, k, k + 1, \dots, n - 2, n - 1$ . Then,  $\det(\mu I - L_c(K_{a_n}(k)))$  will reduce to a new determinant, say  $(\mu - n + 1)^{k-2}(\mu - n)^{n-k-2} \det(C)$ .

Step 2: In  $\det(C)$ , replacing  $C_i$  by  $C'_i = C_i + C_{i-1}$ , for  $i = 4, 5, \dots, k + 1, k + 3, \dots, n$ , a new determinant say  $\det(D)$  is obtained.

Step 3: On expanding  $\det(D)$  along the rows  $R_i$ , for  $i = 3, 4, \dots, k, k + 2, \dots, n - 2, n - 1$ , it reduces to

$$\begin{aligned} \det(D) &= \begin{vmatrix} \mu - n + k + 1 & -1 & 0 & n - k - 1 \\ -1 & \mu - n + 1 & n - \mu - 1 & 0 \\ 0 & 1 & \mu - n + k & n - k - 1 \\ 1 & 1 & k - 1 & \mu - k - 1 \end{vmatrix} \\ &= (\mu^4 - (3n - k - 2)\mu^3 + ((3n^2 - 4n) - (2n - 1)k)\mu^2 - ((n^3 - 2n^2 + 2n - 3) - (n^2 - n + 2)k)\mu + (2n^2 - 6n + 3) - (2n - 3)k). \end{aligned}$$

Thus,

$$\phi(L_c(K_{a_n}(k), \mu)) = (\mu - n)^{n-k-2}(\mu - n + 1)^{k-2}(\mu^4 - (3n - k - 2)\mu^3 + ((3n^2 - 4n) - (2n - 1)k)\mu^2 - ((n^3 - 2n^2 + 2n - 3) - (n^2 - n + 2)k)\mu + (2n^2 - 6n + 3) - (2n - 3)k).$$

□

**Definition 2.6.** For  $n \geq 3$  and  $0 \leq k \leq \lfloor \frac{n}{3} \rfloor$ , the graph  $K_{e_n}(k)$  is obtained from  $K_n$  by the deletion of  $k$  independent paths  $P_3$ .

**Example 2.2.** Consider the cluster graph obtained from  $K_7$  deleting two independent paths  $\{v_1, v_3, v_5\}$  and  $\{v_2, v_4, v_6\}$  respectively. Let  $c(v_1) = c(v_3) = 1$ ,  $c(v_2) = c(v_4) = 3$ ,  $c(v_5) = 2$ ,  $c(v_6) = 4$  and  $c(v_7) = 5$ .




FIGURE 2. Graph  $K_{e_7}(2)$

**Theorem 2.3.** For  $n \geq 3$  and  $1 \leq k \leq \lfloor \frac{n}{3} \rfloor$ ,  $\phi(L_c(K_{e_n}(k), \mu)) = (\mu - n + 1)^{k-1}(\mu - n)^{n-3k-1}(\mu^2 - (2n - 3)\mu + n^2 - 3n - 3)^{k-1}(\mu^4 - (3n - 4)\mu^3 + (3n^2 - 8n + 2k)\mu^2 + (k(10 - 4n) - (n^3 - 4n^2 + 3))\mu + (2n^2 - 10n + 9)k).$

*Proof.* Consider the graph  $K_{e_n}(k)$ . Since  $\chi(K_{e_n}(k)) = n - k$ , we have

$$L_c(K_{e_n}(k)) = \begin{bmatrix} (n-1)I_k - J_k & (-J + 2I)_k & -J_k & -J_{k \times (n-3k)} \\ (-J + 2I)_k & (n-2)I_k - J_k & (-J + I)_k & -J_{k \times (n-3k)} \\ -J_k & (-J + I)_k & (n-1)I_k - J_k & -J_{k \times (n-3k)} \\ -J_{(n-3k) \times k} & -J_{(n-3k) \times k} & -J_{(n-3k) \times k} & (nI - J)_{(n-3k)} \end{bmatrix}$$

Consider  $\det(\mu I - L_c(K_{e_n}(k)))$ .

Step 1: Replace  $R_i$  by  $R'_i = \begin{cases} R_i - R_{i+1}, & \text{for } i=1, 2, \dots, k-1, k+1, k+2, \dots, \\ & 2k-1, 2k+1, 2k+2, \dots, 3k-1, \\ R_i - R_{i-1}, & \text{for } i=n, n-1, \dots, 3k+2. \end{cases}$

Then,  $\det(\mu I - L_c(K_{e_n}(k)))$  will reduce to a new determinant, say  $\det(C)$ .

Step 2: In  $\det(C)$ , replacing

$$C_i \text{ by } C'_i = \begin{cases} C_i + C_{i+1}, & \text{for } i=n-1, n-2, \dots, 3k+1, \\ C_i + C_{i-1} + \dots + C_1, & \text{for } i=k, k-1, \dots, 2, \\ C_i + C_{i-1} + \dots + C_{k+1}, & \text{for } i=2k, 2k-1, \dots, k+2, \\ C_i + C_{i-1} + \dots + C_{2k+1}, & \text{for } i=3k, 3k-1, \dots, 2k+2. \end{cases}$$

a new determinant say  $\det(D)$  is obtained.

Step 3: In  $\det(D)$ , replacing  $C_i$  by  $C'_i = (\mu - n + 1)C_i + 2C_j$ , for  $i = k + 1, k + 2, \dots, 2k - 1$  and  $j = 1, 2, \dots, k - 1$  a new determinant say  $\det(E)$  is obtained.

Step 4: On expanding  $\det(E)$  along the rows  $R_i$ , for  $i = 1, 2, \dots, k-1, 3k+2, 3k+3, \dots, n-1, n$ , it reduces to  $(\mu - n)^{n-3k-1} \det(F)$  which is of order  $2k + 2$ . Where

$$\det(F) = \begin{vmatrix} P & \mu - n + 3 & \dots & N & k - 2 & 1 & \dots & k - 1 & k & n - 3k \\ 0 & M & \dots & 0 & 0 & -1 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & M & 0 & -0 & \dots & -1 & 0 & 0 \\ k - 2 & \mu - n + 3 & \dots & N & \mu - n + k + 2 & 1 & \dots & k - 1 & k - 1 & n - 3k \\ 0 & -\mu + n - 1 & \dots & 0 & 0 & \mu - n + 1 & \dots & 0 & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 0 & 0 & \dots & -\mu + n - 1 & 0 & 0 & \dots & \mu - n + 1 & 0 & 0 \\ k & \mu - n + 3 & \dots & N & k - 1 & 1 & \dots & k - 1 & P & n - 3k \\ k & \mu - n + 3 & \dots & N & k & 1 & \dots & k - 1 & k & \mu - 3k \end{vmatrix}$$

where  $M = \mu^2 - (2n - 3)\mu + n^2 - 3n - 2$ ,  $P = \mu - n - k + 1$  and  $N = (k - 1)(\mu - n + 3)$ .

Step 5: In  $\det(F)$ , replacing  $C_i$  by  $C'_i = C_i + C_{k+i}$ , for  $i = 2, 3, \dots, k - 1$  and then expanding  $\det(E)$  along the rows  $R_i$ , for  $i = 2, 3, \dots, k, k + 2, k + 3, \dots, 2k - 1, 2k$ , it reduces to

$$\det(F) = (\mu - n)^{n-3k-1} (\mu - n + 1)^{k-1} (\mu^2 - (2n - 3)\mu + n^2 - 3n - 3)^{k-1}$$

$$\begin{vmatrix} \mu - n + k + 1 & k - 2 & k & n - 3k \\ k - 2 & \mu - n + k + 2 & k - 1 & n - 3k \\ k & k - 1 & \mu - n + k + 1 & n - 3k \\ k & k & k & \mu - 3k \end{vmatrix}$$

$$= (\mu - n + 1)^{k-1} (\mu^2 - (2n - 3)\mu + n^2 - 3n - 3)^{k-1} (\mu^4 - (3n - 4)\mu^3 + (3n^2 - 8n + 2k)\mu^2 + (k(10 - 4n) - (n^3 - 4n^2 + 3))\mu + (2n^2 - 10n + 9)k).$$

Thus,

$$\begin{aligned} \phi(L_c(K_{e_n}(k), \mu)) &= (\mu - n + 1)^{k-1} (\mu - n)^{n-3k-1} (\mu^2 - (2n - 3)\mu + n^2 - 3n - 3)^{k-1} \\ & (\mu^4 - (3n - 4)\mu^3 + (3n^2 - 8n + 2k)\mu^2 + (k(10 - 4n) - (n^3 - 4n^2 + 3))\mu + \\ & (2n^2 - 10n + 9)k). \end{aligned}$$

□

**Definition 2.7.** For fixed integers  $n \geq 4$  and  $0 \leq k \leq \lfloor \frac{n}{4} \rfloor$ , the graph  $K_{d_n}(k)$  is obtained from  $K_n$  by the deletion of  $k$  independent paths  $P_4$ .

**Example 2.3.** Consider the cluster graph on 7 vertices obtained by deleting one independent path  $\{v_1, v_3, v_4, v_2\}$ . Let  $c(v_1) = c(v_3) = 1$ ,  $c(v_2) = c(v_4) = 2$ ,  $c(v_5) = 3$ ,  $c(v_6) = 4$  and  $c(v_7) = 5$ .




FIGURE 3. Graph  $K_{d_7}(1)$

The color Laplacian matrix is,  $L_c(K_{d_n}(k)) = \left[ \begin{array}{c|c|c} 6I_2 - J_2 & (-J + 2I)_2 & -J_{2 \times 3} \\ \hline (-J + 2I)_2 & 4I_2 & -J_{2 \times 3} \\ \hline -J_{3 \times 2} & -J_{3 \times 2} & (7I - J)_3 \end{array} \right]$

**Theorem 2.4.** For  $n \geq 4$  and  $1 \leq k \leq \lfloor \frac{n}{4} \rfloor$ ,

$$LE_c(K_{d_n}(k)) = \frac{1}{n} \left[ 2k((3 + \sqrt{5})n - 6) - 24k^2 + n(n - 3 + \sqrt{(n + 1)^2 - 16k}) \right].$$

*Proof.* Consider the cluster graph  $K_{d_n}(k)$ . As  $\chi(K_{d_n}(k)) = n - 2k$ , we have

$$L_c(K_{d_n}(k)) = \left[ \begin{array}{c|c|c} (n - 1)I_{2k} - J_{2k} & (-J + 2I)_{2k} & -J_{2k \times (n - 4k)} \\ \hline (-J + 2I)_{2k} & X_{2k} & -J_{2k \times (n - 4k)} \\ \hline -J_{(n - 4k) \times 2k} & -J_{(n - 4k) \times 2k} & (nI - J)_{(n - 4k)} \end{array} \right]$$

$$X = \left[ \begin{array}{c|c|c|c} (n - 3)I_{2 \times 2} & -J_{2 \times 2} & \cdots & -J_{2 \times 2} \\ \hline -J_{2 \times 2} & (n - 3)I_{2 \times 2} & \cdots & -J_{2 \times 2} \\ \hline \vdots & \vdots & \ddots & \vdots \\ \hline -J_{2 \times 2} & -J_{2 \times 2} & \cdots & (n - 3)I_{2 \times 2} \end{array} \right]$$

Consider  $\det(\mu I - L_c(K_{d_n}(k)))$ .

Step 1: Replace  $R_i$  by  $R'_i = \begin{cases} R_i - R_{i+1}, & \text{for } i = 1, 2, \dots, 2k-1, 2k+1, \\ R_i - R_{i-1}, & \text{for } i = n, n-1, \dots, 4k+2. \end{cases}$

Then,  $\det(\mu I - L_c(K_{d_n}(k)))$  will reduce to a new determinant, say  $\det(C)$ .

Step 2: In  $\det(C)$ , replacing

$$C_i \text{ by } C'_i = \begin{cases} C_i + C_{i+1}, & \text{for } i = n-1, n-2, \dots, 4k+2, 4k+1, \\ C_i + C_{i-1} + \dots + C_1, & \text{for } i = 2k, 2k-1, \dots, 2, \\ C_i + C_{i-1} + \dots + C_{2k}, & \text{for } i = 4k, 4k-1, \dots, 2k+2. \end{cases}$$

a new determinant say  $\det(D)$  is obtained.

Step 3: In  $\det(D)$ , replace  $C_i$  by  $C'_i = (\mu - n + 1)C_i + 2C_j$ , for  $i = 2k + 1, 2k + 2, \dots, 4k$  and  $j = 1, 2, \dots, 2k - 1$ . It reduces to  $\det(E)$ .

Step 4: On expanding  $\det(E)$  along the rows  $R_i$ , for  $i = 1, 2, \dots, 2k - 1, 2k + 1, 2k + 3, \dots, 4k - 3, 4k - 1, 4k + 2, 4k + 3, \dots, n - 1, n$ , it becomes  $(\mu - n)^{n-4k+1}(\mu^2 - (2n - 4)\mu + n^2 - 4n - 1)^k \det(F)$ . Where

$$\det(F) = \begin{vmatrix} P & 2(\mu - n + 3) & 4(\mu - n + 3) & \cdots & (2k - 2)(\mu - n + 3) & 2k - 2 & n - 4k \\ 2k - 2 & Q & R & \cdots & S & P & n - 4k \\ 2k - 2 & 2(\mu - n + 3) & R & \cdots & S & P & n - 4k \\ 2k - 2 & 2(\mu - n + 3) & 4(\mu - n + 3) & \cdots & S & P & n - 4k \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots \\ 2k - 2 & 2(\mu - n + 3) & 4(\mu - n + 3) & \cdots & S & P & n - 4k \\ 2k - 2 & 2(\mu - n + 3) & 4(\mu - n + 3) & \cdots & (2k - 2)(\mu - n + 3) & P & n - 4k \\ 2k - 2 & 2(\mu - n + 3) & 4(\mu - n + 3) & \cdots & (2k - 2)(\mu - n + 3) & 2k & \mu - 4k \end{vmatrix}$$

where  $P = \mu - n + 2k + 1$ ,  $Q = \mu^2 + 2\mu(2 - n) + n^2 - 4n + 3$ ,  $R = \mu^2 + 2\mu(3 - n) + n^2 - 6n + 9$  and  $S = \mu^2 + 2\mu(k - n) + n^2 - k(2n - 6) - 9$

Step 5: In  $\det(F)$ , replacing  $R_i$  by  $R'_i = R_i - R_{i+1}$ , for  $i = 2, 3, \dots, k, k + 1$ , it reduces to

$$\det(F) = (\mu^2 - (n - 1)2\mu + n^2 - 2n - 3)^{k-1} \begin{vmatrix} \mu - n + 2k + 1 & 2k - 2 & n - 4k \\ -2 & \mu - n + 1 & n - \mu \\ 2k & 2k & \mu - 4k \end{vmatrix}$$



$$= (\mu^2 - (n - 1)2\mu + n(n - 2) - 3)^{k-1}(\mu^3 + (2 - 2n)\mu^2 - (n^2 - 2n + 4k - 3)\mu + 4k(3 - n)).$$

Hence, by back substitution, we obtain characteristic polynomial of cluster graph  $K_{d_n}(k)$ ,

$$\phi(L_c(K_{d_n}(k), \mu)) = (\mu - n + 3)^k(\mu - n - 1)^{k-1}(\mu - n)^{n-4k-1}(\mu^2 - (n + 1)\mu + 4k)(\mu^2 - (2n - 4)\mu + (n^2 - 4n - 1))^k.$$

So, the color Laplacian spectrum of  $K_{d_n}(k)$  is  $\{n - 3$  ( $k$  times),  $n + 1$  ( $k - 1$  times),  $n(n - 4k - 1)$  times,  $n - 2 + \sqrt{5}$  ( $k$  times),  $n - 2 - \sqrt{5}$  ( $k$  times),  $\frac{n + 1 + \sqrt{(n + 1)^2 - 16k}}{2}$ ,  $\frac{n + 1 - \sqrt{(n + 1)^2 - 16k}}{2}\}$ .

Average degree of  $K_{d_n}(k)$  is  $\frac{n^2 - n - 6k}{n}$ . Hence,

$$\begin{aligned} LE_c(K_{d_n}(k)) &= k \left| n - 3 - \frac{n^2 - n - 6k}{n} \right| + (k - 1) \left| n + 1 - \frac{n^2 - n - 6k}{n} \right| + \\ & (n - 4k - 1) \left| n - \frac{n^2 - n - 6k}{n} \right| + \left| \frac{n + 1 \pm \sqrt{(n + 1)^2 - 16k}}{2} - \frac{n^2 - n - 6k}{n} \right| + k \left| n - 2 \pm \sqrt{5} - \frac{n^2 - n - 6k}{n} \right| \\ &= \frac{1}{n} \left[ 2k((3 + \sqrt{5})n - 6) - 24k^2 + n(n - 3 + \sqrt{(n + 1)^2 - 16k}) \right]. \end{aligned}$$

□

### 3. COLOR LAPLACIAN ENERGY OF BIPARTITE CLUSTER GRAPHS

**Definition 3.1.** [14] Let  $K_{m,n}$  be the complete bipartite graph,  $1 \leq r \leq m$ ,  $1 \leq s \leq n$  and  $m, n \geq 1$  be the integers. The bi-cluster graph  $Kb_{m,n}(r, s)$  is obtained by deleting the edges of  $K_{r,s}$  from  $K_{m,n}$ .

**Example 3.1.** Consider the bi-cluster graph. Let  $c(v_1) = c(v_2) = c(v_3) = c(v_4) = 1$  and  $c(v_5) = c(v_6) = c(v_7) = c(v_8) = 2$ .




FIGURE 4. Graph  $Kb_{4,4}(2, 2)$

**Theorem 3.1.** For  $m, n \geq 1$  and  $0 \leq r \leq m$ ,  $0 \leq s \leq n$ ,

$$\begin{aligned} \phi(L_c(Kb_{m,n}(r,s),\mu)) &= (\mu - n + s + 1)^{r-1}(\mu - m + r + 1)^{s-1}(\mu - n + 1)^{m-r-1}(\mu - \\ & m + 1)^{n-s-1}(\mu^4 - (3(n+m) - r - s - 4)\mu^3 + (3n^2 + (6m - 9)n + 3(m^2 - 3m + 2) - \\ & (3n + 2m - 3)r - (2n + 3m - 4r - 3)s)\mu^2 - (n^3 + (4m - 6)n^2 + (4m^2 - 12m + 9)n \\ & + m^3 - 6m^2 + 9m - 4 - (3n^2 + (3m - 6)n + m^2 - 4m + 3)r - (n^2 + (3m - 4)n + \\ & 3m^2 - 6m + 3 - (4r(n + m - 2))s)\mu + ((m - 1)n^3 + (m^2 - 4m + 3)n^2 + (m^3 - 4m^2 \\ & + 6m - 3)n - (m^3 - 3m^2 + 3m - 1) - (n^3 + (m - 3)n^2 + (m^2 - 3m + 3)n - m^2 + \\ & 2m - 1)r - ((m - 1)n^2 + (m^2 - 3m + 2)n + (m^3 - 3m^2 + 3m - 1) - (n^2 + (2m - 4) \\ & n + (m^2 - 4m + 4))r)s). \end{aligned}$$

*Proof.* Let  $U = \{u_1, u_2, \dots, u_r, u_{r+1}, \dots, u_m\}$  and

$V = \{v_1, v_2, \dots, v_s, v_{s+1}, \dots, v_n\}$  be the partites of complete bipartite graph  $K_{m,n}$ . The graph  $Kb_{m,n}(r,s)$  is obtained by deleting the edges of  $K_{r,s}$  from  $K_{m,n}$ . Note that,  $\chi(Kb_{m,n}(r,s)) = 2$ . Then, we have

$$L_c(Kb_{m,n}(r,s)) = \begin{bmatrix} J_r + (n-s-1)I_r & J_{r \times (m-r)} & 0_{r \times s} & -J_{r \times (n-s)} \\ J_{(m-r) \times r} & J_r + (n-1)I_{(m-r)} & -J_{(m-r) \times s} & -J_{(m-r) \times (n-s)} \\ 0_{s \times r} & -J_{s \times (m-r)} & J_r + (m-r-1)I_s & J_{s \times (n-s)} \\ -J_{n-s \times r} & -J_{(n-s) \times (m-r)} & J_{(n-s) \times s} & (J + (m-1)I)_{(n-s)} \end{bmatrix}$$

Consider  $\det(\mu I - L_c(Kb_{m,n}(r,s)))$ .

Step 1: Replace  $R_i$  by  $R'_i = R_i - R_{i+1}$ , for  $i = 1, 2, \dots, r-1, r+1, r+2, \dots, m-1, 1, 2, \dots, s-1, s+1, s+2, \dots, n-1$ . Then,  $\det(\mu I - L_c(Kb_{m,n}(r,s)))$  will reduce to a new determinant, say  $(\mu - n + s + 1)^{r-1}(\mu - m + r + 1)^{s-1}(\mu - n + 1)^{m-r-1}(\mu - m + 1)^{n-s-1} \det(C)$ .

Step 2: In  $\det(C)$ , replacing

$$C_i \text{ by } C'_i = \begin{cases} C_i + C_{i-1} + \dots + C_1, & \text{for } i=r, r-1, \dots, 2 \\ C_i + C_{i-1} + \dots + C_1, & \text{for } i=s, s-1, \dots, 2, \\ C_i + C_{i-1} + \dots + C_{r+1}, & \text{for } i=m, m-1, \dots, r+2, \\ C_i + C_{i-1} + \dots + C_{s+1}, & \text{for } i=n, n-1, \dots, s+2. \end{cases}$$

a new determinant say  $\det(D)$  is obtained.

Step 3: On expanding  $\det(D)$  along the rows  $R_i$ , for  $i = 1, 2, \dots, r-1, r+1, r+2, \dots, m-1, 1, 2, \dots, s-1, s+1, s+2, \dots, n-1$ , it reduces to

$$\det(D) = \begin{vmatrix} \mu - n + s - r + 1 & r - m & 0 & n - s \\ -r & \mu - m - n + r + 1 & s & n - s \\ 0 & m - r & \mu - m + r - s + 1 & s - n \\ r & m - r & -s & \mu - m - n + s + 1 \end{vmatrix}$$

$$\begin{aligned} &= (\mu^4 - (3(n+m) - r - s - 4)\mu^3 + (3n^2 + (6m - 9)n + 3(m^2 - 3m + 2) - (3n + \\ & 2m - 3)r - (2n + 3m - 4r - 3)s)\mu^2 - (n^3 + (4m - 6)n^2 + (4m^2 - 12m + 9)n + m^3 - \\ & 6m^2 + 9m - 4 - (3n^2 + (3m - 6)n + m^2 - 4m + 3)r - (n^2 + (3m - 4)n + 3m^2 - 6m + \\ & 3 - (4r(n + m - 2))s)\mu + ((m - 1)n^3 + (m^2 - 4m + 3)n^2 + (m^3 - 4m^2 + 6m - 3)n - \\ & (m^3 - 3m^2 + 3m - 1) - (n^3 + (m - 3)n^2 + (m^2 - 3m + 3)n - m^2 + 2m - 1)r - ((m - \\ & 1)n^2 + (m^2 - 3m + 2)n + (m^3 - 3m^2 + 3m - 1) - (n^2 + (2m - 4)n + (m^2 - 4m + 4))r)s). \end{aligned}$$

Thus,

$$\begin{aligned} \phi(L_c(Kb_{m,n}(r,s),\mu)) &= (\mu - n + s + 1)^{r-1}(\mu - m + r + 1)^{s-1}(\mu - n + 1)^{m-r-1}(\mu - \\ & m + 1)^{n-s-1}(\mu^4 - (3(n+m) - r - s - 4)\mu^3 + (3n^2 + (6m - 9)n + 3(m^2 - 3m + 2) - \\ & (3n + 2m - 3)r - (2n + 3m - 4r - 3)s)\mu^2 - (n^3 + (4m - 6)n^2 + (4m^2 - 12m + 9)n \\ & + m^3 - 6m^2 + 9m - 4 - (3n^2 + (3m - 6)n + m^2 - 4m + 3)r - (n^2 + (3m - 4)n + \\ & 3m^2 - 6m + 3 - (4r(n + m - 2))s)\mu + ((m - 1)n^3 + (m^2 - 4m + 3)n^2 + (m^3 - 4m^2 \\ & + 6m - 3)n - (m^3 - 3m^2 + 3m - 1) - (n^3 + (m - 3)n^2 + (m^2 - 3m + 3)n - m^2 + \\ & 2m - 1)r - ((m - 1)n^2 + (m^2 - 3m + 2)n + (m^3 - 3m^2 + 3m - 1) - (n^2 + (2m - 4) \\ & n + (m^2 - 4m + 4))r)s) \end{aligned}$$

□

**Definition 3.2.** [14] Let  $e_i, i = 1, 2, \dots, k, 1 \leq k \leq \min\{m, n\}$ , be independent edges of the complete bipartite graph  $K_{m,n}, m, n \geq 1$ . The graph  $Ka_{m,n}(k)$  is obtained by deleting  $e_i, i = 1, 2, \dots, k$  from  $K_{m,n}$ .

**Theorem 3.2.** For  $m, n \geq 1$  and  $0 \leq k \leq \min\{m, n\}$

$$\begin{aligned} \phi(L_c(Ka_{m,n}(k),\mu)) &= (\mu - m + 1)^{n-k-1}(\mu - n + 1)^{m-k-1}(\mu^2 - (n + m - 4)\mu + \\ & (n(m - 2) - (2m - 3)))^{k-1}(\mu^4 - 3(m + n - 2)\mu^3 + (3(n^2 + m^2) + (6m - 14)n - \\ & 14m + 4k + 12)\mu^2 - (n^3 + (4m - 10)n^2 + (4m^2 - 18m + 19)n + 4k(n + m - 2) \\ & + m^3 - 10m^2 - 19m - 10)\mu + ((m - 2)n^3 + (m^2 - 6m + 7)n^2 + (m^3 - 6m^2 + \\ & 12m - 8)n - (2m^3 - 7m^2 + 8m - 3) + (n^2 + (2m - 4)n + (m^2 - 4m + 4))k). \end{aligned}$$

*Proof.* Let  $U = \{u_1, u_2, \dots, u_k, u_{k+1}, \dots, u_m\}$  and

$V = \{v_1, v_2, \dots, v_k, v_{k+1}, \dots, v_n\}$  be the partites of complete bipartite graph  $K_{m,n}$ . The graph  $Ka_{m,n}(k)$  is obtained by deleting independent edges  $e_i$  of the complete bipartite graph  $K_{m,n}, i = 1, 2, \dots, k$ . Observe that  $\chi(Ka_{m,n}(k)) = 2$ . Then, we have

$$L_c(Ka_{m,n}(k)) = \begin{bmatrix} J_k + (n - 2)I_k & J_{k \times (m-k)} & (-J + I)_k & -J_{k \times (n-k)} \\ J_{(m-k) \times k} & (J + (n - 1)I)_{(m-k)} & -J_{(m-k) \times k} & -J_{(m-k)} \\ (-J + I)_k & -J_{k \times (m-k)} & J_k + (m - 2)I_k & J_{k \times (n-k)} \\ -J_{(n-k) \times k} & -J_{(n-k) \times (m-k)} & J_{(n-k) \times k} & (J + (m - 1)I)_{(n-k)} \end{bmatrix}$$

Consider  $\det(\mu I - L_c(Ka_{m,n}(k)))$ .

Step 1: Replace  $R_i$  by  $R'_i = R_i - R_{i+1}$ , for  $i = u_1, u_2, \dots, u_{k-1}, u_{k+1}, u_{k+2}, \dots, u_{m-1}, v_1, v_2, \dots, v_{k-1}, v_{k+1}, v_{k+2}, \dots, v_{n-1}$ . Then,  $\det(\mu I - L_c(Ka_{m,n}(k)))$  will reduce to a new determinant, say  $(\mu - n + 1)^{m-k-1}(\mu - m + 1)^{n-k-1} \det(C)$ .

Step 2: In  $\det(C)$ , replacing

$$C_i \text{ by } C'_i = \begin{cases} C_i + C_{i-1} + \dots + C_1, & \text{for } i = u_k, u_{k-1}, \dots, u_2, \\ C_i + C_{i-1} + \dots + C_1, & \text{for } i = v_k, u_{k-1}, \dots, v_2, \\ C_i + C_{i-1} + \dots + C_{k+1}, & \text{for } i = u_m, u_{m-1}, \dots, u_{k+2}, \\ C_i + C_{i-1} + \dots + C_{k+1}, & \text{for } i = v_n, u_{n-1}, \dots, v_{k+2}. \end{cases}$$

a new determinant say  $\det(D)$  is obtained.

Step 3: In  $\det(D)$ , replacing  $C_i$  by  $C'_i = (\mu - n + 2)C_i + C_j$ , for  $i = v_1, v_2, \dots, v_{k-1}$  and  $j = u_1, u_2, \dots, u_{k-1}$ , it simplifies to a determinant say  $\det(E)$ .

Step 4: On expanding  $\det(E)$  along the rows  $R_i$ , for  $i = u_1, u_2, \dots, u_{k-1}, u_{k+1}, u_{k+2}, \dots, u_{m-1}, v_1, v_2, \dots, v_{k-1}, v_{k+1}, v_{k+2}, \dots, v_{n-1}$ , it reduces to

$$\det(F) = (\mu^2 - (n + m - 4)\mu + (n(m - 2) - (2m - 3)))^{k-1}$$

$$\begin{vmatrix} \mu - n - k + 2 & k - m & k - 1 & n - k \\ -k & \mu - n - m + k + 1 & k & n - k \\ k - 1 & m - k & \mu - m - k + 2 & k - n \\ k & m - k & -k & \mu - m - n + k + 1 \end{vmatrix}$$

$$= (\mu^2 - (n + m - 4)\mu + (n(m - 2) - (2m - 3)))^{k-1} (\mu^4 - 3(m + n - 2)\mu^3 + (3(n^2 + m^2) + (6m - 14)n - 14m + 4k + 12)\mu^2 - (n^3 + (4m - 10)n^2 + (4m^2 - 18m + 19)n + 4k(n + m - 2) + m^3 - 10m^2 - 19m - 10)\mu + ((m - 2)n^3 + (m^2 - 6m + 7)n^2 + (m^3 - 6m^2 + 12m - 8)n - (2m^3 - 7m^2 + 8m - 3) + (n^2 + (2m - 4)n + (m^2 - 4m + 4))k)).$$

Thus,

$$\begin{aligned} \phi(L_c(Ka_{m,n}(k), \mu)) &= (\mu - m + 1)^{n-k-1} (\mu - n + 1)^{m-k-1} (\mu^2 - (n + m - 4)\mu + \\ & (n(m - 2) - (2m - 3)))^{k-1} (\mu^4 - 3(m + n - 2)\mu^3 + (3(n^2 + m^2) + (6m - 14)n - \\ & 14m + 4k + 12)\mu^2 - (n^3 + (4m - 10)n^2 + (4m^2 - 18m + 19)n + 4k(n + m - 2) + \\ & m^3 - 10m^2 - 19m - 10)\mu + ((m - 2)n^3 + (m^2 - 6m + 7)n^2 + (m^3 - 6m^2 + 12m \\ & - 8)n - (2m^3 - 7m^2 + 8m - 3) + (n^2 + (2m - 4)n + (m^2 - 4m + 4))k)). \end{aligned}$$

□

#### 4. CONCLUSION

The energy, color energy and color Laplacian energy of a graph are the emerging concepts within graph theory. In this paper we have evaluated color Laplacian energy of Cluster and bi-cluster graphs. The color Laplacian spectrum of cluster and bi-cluster graphs is expressed in terms of its parameters.

#### REFERENCES

- [1] Abreu, N. M. M., Vinagre, C. T. M., Bonifacio, A. S. and Gutman, I., (2008), The Laplacian energy of some Laplacian integral graphs, MATCH Commun. Math. Comput. Chem., 60, pp. 447-460.
- [2] Adiga, C., Sampathkumar, E., Sriraj, M. A. and Shrikanth, A. S., (2013), Color energy of a graph, Proc. Jangjeon Math. Soc., 16, pp. 335-351.
- [3] Adiga, C., Sampathkumar, E. and Sriraj, M. A., (2014), Color energy of unitary cayley graphs, Discussiones Mathematicae Graph Theory, 34, pp. 707-721.
- [4] Balakrishnan, R. (2004), The energy of a graph, Linear Algebra Appl., 387, pp. 287-295.
- [5] Bapat, R. B., (2011), Graphs and matrices, Springer-Hindustan Book Agency, London.
- [6] Bapat, R. B. and Pati, S., (2004), Energy of a graph is never an odd integer, Bulletin of Kerala Mathematical Association, 1, pp. 129-132.
- [7] Bhat, P. G. and D'Souza, S., (2015), Color Laplacian energy of a graph, Proc. Jangjeon Math. Soc., 18, pp. 321-330.
- [8] Bhat, P. G. and D'Souza, S., (2017), Color signless Laplacian energy of graphs, AKCE International Journal of Graphs and Combinatorics, 14, pp. 142-148.
- [9] Gutman, I., (1978), The energy of a graph, Ber. Math. Stat. Sect. Forschungsz. Graz., 103, pp. 1-22.
- [10] Gutman, I. and Zhou, B., (2006), Laplacian energy of a graph, Linear Algebra Appl., 414, pp. 29-37.
- [11] Gutman, I. and Pavlovic, L., (1999), The energy of some graphs with large number of edges, Bull. Acad. Serbe Sci. Arts (Cl. Math. Natur.), 118, pp. 35-50.
- [12] Harary, F., (1989), Graph Theory, Narosa Publishing House, New Delhi.
- [13] Walikar, H. B. and Ramane, H. S., (2001), Energy of some cluster graphs, Kragujevac J. Sci., 23, pp. 51-62.
- [14] Walikar, H. B. and Ramane, H. S., (2001), Energy of some bipartite cluster graphs, Kragujevac J. Sci., 23, pp. 63-74.



**Sabitha D'Souza** received her B.Sc. and M.Sc degrees in Mathematics from Mangalore University, India, in 2001 and 2003, respectively. She received her Ph.D. from Manipal Academy of Higher Education, Manipal, India in 2016. She is currently working as an assistant professor at Manipal Institute of Technology, Manipal Academy of Higher Education, Manipal. Her research interests include Graph coloring, graph complements and Spectral graph theory.

---



**Gowtham H. J.** received his B.Sc. degree from Mangalore University, Mangalore, India, in 2012. He received his M.Sc. degree from Manipal Academy of Higher Education, Manipal, in 2014. He received his Ph.D. from Manipal Academy of Higher Education, in 2020. He is currently working as an assistant professor at Manipal Institute of Technology, Manipal Academy of Higher Education. His research interests include Spectral graph theory and Graph Labeling.

---



**Pradeep G. Bhat** received his B.Sc. and M. Sc degrees in Mathematics from Karnataka University, Dharawad, India, in 1984 and 1986, respectively. He received his Ph.D. from Mangalore University, in 1998. At present, he is a professor in the Department of Mathematics at Manipal Institute of Technology, Manipal Academy of Higher Education. He served as a HOD of Mathematics from 2013 to 2018 at MIT, Manipal. His research interests include Graph complements, Spectral graph theory and Graph labeling.

---



**Girija K. P.** received her B.Sc. Degree from Kuvempu University, Shimoga, India, in 2013. She received her M. Sc. degree from Kuvempu University, Shimoga, India in 2015. At present, she is pursuing her Ph.D. under the guidance of Dr. Pradeep G. Bhat, Dr. Devadas Nayak C. and Dr. Sabitha D'Souza at Manipal Institute of Technology, Manipal Academy of Higher Education, Manipal. Her research interests include Spectral graph theory and Graph Labeling.

---

---