

SOME PROPERTIES OF ZERO-DIVISOR GRAPHS OF DIRECT PRODUCTS OF FINITE FIELDS

SUBHASH MALLINATH GADED^{1*}, NITHYA SAI NARAYANA², §

ABSTRACT. This study investigates the zero-divisor graphs of the direct products of finite fields with vertex set consisting of non-zero zero divisors of the direct products of finite fields, and two distinct non-zero zero divisors are adjacent in the zero divisor graph if the product of the two distinct non-zero zero divisors is the additive identity of the direct products of finite fields. We prove that the metric chromatic number, clique number, vertex chromatic number are all equal to n , for the zero divisor graph of direct products of n finite fields, and also find the metric chromatic number, clique number, and vertex chromatic number of the complement graph of the zero-divisor graph of the direct product of n fields. The independence number, edge chromatic number, Eulerian and Hamiltonian properties of the zero-divisor graph and the complement graph are also determined.

Keywords: Zero-divisor graph, complement graph, finite field.

AMS Subject Classification: 05C15, 05C25, 05C45.

1. INTRODUCTION

In 1988, I. Beck [1] defined a graph on a commutative ring R , by considering all the elements of the ring R as the vertices of the graph and two distinct vertices x and y are adjacent if and only if $x \cdot y = 0$. In 1999, this definition of the graph given by Beck was modified by Anderson and Livingston [2] in which they considered only the non-zero zero divisors of R to be the vertex set of the zero divisor graph of R denoted by $\Gamma(R)$ and two distinct vertices x and y are adjacent in $\Gamma(R)$ if and only if $x \cdot y = 0$. In this paper we consider the zero divisor graph of the ring, $F_1 \times \cdots \times F_n$, ($n \geq 2$), where $F_1, \dots, F_n$ are finite fields.

Let $Z^*(F_1 \times \cdots \times F_n)$ be the set of non-zero zero-divisors of the ring $F_1 \times \cdots \times F_n$, ($n \geq 2$) and $\Gamma(F_1 \times \cdots \times F_n)$ (defined by Anderson and Livingston [2]) denote the graph with vertex set $Z^*(F_1 \times \cdots \times F_n)$. Let $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n) \in V(\Gamma(F_1 \times \cdots \times F_n)) = Z^*(F_1 \times \cdots \times F_n)$. Note that at least one coordinate of x and at least one coordinate of y

¹ R K Talreja College of Arts, Science and Commerce, Ulhasnagar-03, District Thane, India
e-mail: gadedsubhash@gmail.com; ORCID: <https://orcid.org/0000-0002-0883-7273>.

² Department of Mathematics, University of Mumbai, India.
e-mail: narayana_nithya@yahoo.com; ORCID: <https://orcid.org/0000-0001-9335-522X>.

§ Manuscript received: July 20, 2024; accepted: December 07, 2024.

TWMS Journal of Applied and Engineering Mathematics, Vol.15, No.8; © Işık University, Department of Mathematics, 2025; all rights reserved.

contains 0 and at least one coordinate of x and at least one coordinate of y contains non-zero entry. Therefore, the product $x \cdot y = (x_1, \dots, x_n) \cdot (y_1, \dots, y_n) = (x_1 \cdot y_1, \dots, x_n \cdot y_n)$, which is coordinate wise multiplication, is either equal to $(0, \dots, 0)$ or the product $x \cdot y$ contains 0 in at least one coordinate and non-zero entry in at least one coordinate. In other words, $x \cdot y = (0, \dots, 0)$ or $x \cdot y \in Z^*(F_1 \times \dots \times F_n)$. Therefore, the edge set of $\Gamma(F_1 \times \dots \times F_n)$ is

$$\left\{ xy \mid x \cdot y = (0, \dots, 0), x, y \in Z^*(F_1 \times \dots \times F_n) \right\}$$

and the edge set of $\overline{\Gamma(F_1 \times \dots \times F_n)}$ is

$$\left\{ xy \mid x \cdot y \neq (0, \dots, 0), x, y \in Z^*(F_1 \times \dots \times F_n) \right\}.$$

The complement graph [5] $\overline{G}$ of a simple graph G is defined by taking $V(\overline{G}) = V(G)$ and making two vertices u and v adjacent in $\overline{G}$ if and only if they are nonadjacent in G . Thus, if $x \cdot y = (0, \dots, 0)$, then $xy \in E(\Gamma(F_1 \times \dots \times F_n))$, and if $x \cdot y \neq (0, \dots, 0)$, then $xy \in E(\overline{\Gamma(F_1 \times \dots \times F_n)})$ which implies $x \cdot y \in Z^*(F_1 \times \dots \times F_n)$.

1.1. Prerequisites. For basic algebraic terminologies we refer to [3, 4]. For basic graph theoretical terminologies we adopt the definitions of [5, 6, 7].

- (i) The degree of a vertex v in a graph G is the number of edges incident with v and is denoted by $\deg_G(v)$ or simply by $\deg v$. Also, $\deg v$ is the number of vertices adjacent to v . Two adjacent vertices are referred to as neighbors of each other. The set $N(v)$ of neighbors of a vertex v is called the neighborhood of v . Thus $\deg v = |N(v)|$. [6] The minimum degree of G is the minimum degree among the vertices of G and is denoted by $\delta(G)$, the maximum degree of G is the maximum degree among the vertices of G and is denoted by $\Delta(G)$. [6]
- (ii) If every two distinct vertices in a subset $C \subseteq V(G)$ are adjacent in a graph G , the subset is said to be a clique in G . If there isn't another clique in the graph G with more vertices, then a clique C is said to be the maximum clique. The size of the maximal clique in a graph G is its clique number, which is represented by the symbol $\omega(G)$. [6] A set of vertices S in a graph G that has no two adjacent vertices is said to be an independent set. The highest possible size of an independent set in a graph is known as its independence number and is denoted by $\alpha(G)$. [6] A vertex cover of a graph is a collection of vertices such that each edge of the graph is incident to at least one vertex of the set. The cardinality of the smallest such set is known as the minimal vertex cover number and is represented by $\tau(G)$. [6, 7]
- (iii) A graph's k -vertex(edge) coloring is the process of applying k different colors to its vertices(edges) while ensuring that no two adjacent vertices(edges) receive the same color. The least number of colors needed to color a graph's vertices(edges) so that no two adjacent vertices(edges) have the same color is the graph's vertex chromatic number(edge chromatic number), indicated by the symbol $\chi(G)(\chi_1(G))$. [6, 9] A simple graph G is Class-1, if $\chi_1(G) = \Delta(G)$ and Class-2, if $\chi_1(G) = \Delta(G) + 1$. [8]
- (iv) An Eulerian trail is a path through a graph that includes every edge exactly once. It could be open or closed. If a graph has a closed Eulerian trail, it is referred to as an Eulerian graph. [9] A Hamiltonian cycle is a graph cycle that contains every vertex of the graph. A graph is referred to as a Hamiltonian graph if a Hamiltonian cycle exists in it. [9]

1.2. Commonly used results. The following results from graph theory are used to determine the properties of zero divisor graphs of direct product of finite fields.

- (i) For every graph G , $\chi(G) \geq \omega(G)$. [7]
- (ii) For any graph G , $|V(G)| = \alpha(G) + \tau(G)$. [5, 7]
- (iii) If graph G is bipartite, then $\chi_1(G) = \Delta(G)$. [7]
- (iv) If G is simple graph, then $\Delta(G) \leq \chi_1(G) \leq \Delta(G) + 1$. [7]
- (v) If G has a Hamiltonian cycle, then for each non-empty subset $S \subset V$, the subgraph $G - S$ has at most $|S|$ components. [7, 9]
- (vi) If G is a simple graph with at least three vertices and $\delta(G) \geq \frac{n(G)}{2}$, then G is Hamiltonian. [7]
- (vii) A connected graph G is Eulerian if and only if every vertex of G has even degree. [9]

1.3. Known results of zero divisor graphs of direct product of finite fields.

- (i) $|Z^*(F_1 \times \cdots \times F_n)| = |V(\Gamma(F_1 \times \cdots \times F_n))|$
 $= \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1$. [10, Theorem 2.1]
- (ii) If $|F_1| = \cdots = |F_n| = p$, then the number of vertices of degree $p^r - 1$, $(1 \leq r \leq n-1)$ in $\Gamma(F_1 \times \cdots \times F_n)$ is $C(n, r)(p-1)^{n-r}$. [10, Theorem 2.2]
- (iii) If $|F_1| = \cdots = |F_n| = p$, then the number of vertices of degree $p^n - (p-1)^n - p^r - 1$, $(1 \leq r \leq n-1)$ in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is $C(n, r)(p-1)^r$. [10, Corollary 2.3]
- (iv) $\Gamma(F_1 \times F_2)$ is a complete bipartite graph, and hence $\Gamma(F_1 \times F_2)$ is disconnected. [10, Corollary 2.5]

1.4. Observation.

Observation 1. The number of vertices in $\Gamma(F_1 \times \cdots \times F_n)$ with degree

$$\prod_{i_k=1}^r |F_{i_k}| - 1 \text{ is } \frac{\prod_{i=1}^n (|F_i| - 1)}{\prod_{i_k=1}^r (|F_{i_k}| - 1)} \text{ where } 1 \leq i_1 < i_2 < \cdots < i_r \leq n.$$


Observation 2. Let $S = \{v_i = (0, \dots, 0, 1, 0, \dots, 0) : 1 \leq i \leq n\}$. In other words, if $v_i \in S$, then v_i contains 1 in i^{th} coordinate and 0's in remaining coordinates. Clearly, $v_i \cdot v_j = (0, \dots, 0)$ if $i \neq j$, $(1 \leq i \neq j \leq n)$. Therefore, the sub-graph spanned by S is complete graph K_n in $\Gamma(F_1 \times \cdots \times F_n)$.

1.5. Main Results. Let $F_1, \dots, F_n$ ($n \geq 2$) be finite fields. In this paper, we consider the zero-divisor graph $\Gamma(F_1 \times \cdots \times F_n)$ defined by Anderson and Livingston [2], with vertex set $Z^*(F_1 \times \cdots \times F_n)$, and two distinct vertices $x = (x_1, \dots, x_n)$, $y = (y_1, \dots, y_n) \in Z^*(F_1 \times \cdots \times F_n)$ are adjacent in $\Gamma(F_1 \times \cdots \times F_n)$ if and only if $x \cdot y = (0, \dots, 0)$. We prove that the vertex chromatic number $\chi(\Gamma(F_1 \times \cdots \times F_n))$, clique number $\omega(\Gamma(F_1 \times \cdots \times F_n))$, metric chromatic number $\mu(\Gamma(F_1 \times \cdots \times F_n))$ of the zero divisor graph $\Gamma(F_1 \times \cdots \times F_n)$ is n and also determine the vertex chromatic number, clique number, metric chromatic number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. We determine the Edge chromatic number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$, and also determine the independence number, Eulerian and Hamiltonian properties of the zero divisor graph $\Gamma(F_1 \times \cdots \times F_n)$ and the complement graph $\overline{\Gamma(F_1 \times \cdots \times F_n)}$.

2. GRAPH PARAMETERS OF ZERO DIVISOR GRAPH

Example 2.1. Consider the zero divisor graph $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ (as defined in [2]).

$$V(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = \left\{ v_1 = (1, 0, 0), v_2 = (0, 1, 0), v_3 = (0, 0, 1), v_4 = (0, 0, 2), v_5 = (1, 1, 0), \right. \\ \left. v_6 = (1, 0, 1), v_7 = (1, 0, 2), v_8 = (0, 1, 1), v_9 = (0, 1, 2) \right\}.$$


 FIGURE 1. $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$

- (i) Consider the subset $S = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\} \subset V(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3))$. The subgraph induced by S is complete graph K_3 .
- (ii) Consider the sub-partition $V_1 = \{(1, 0, 0), (1, 0, 1), (1, 0, 2), (1, 1, 0)\}$, $V_2 = \{(0, 1, 0), (0, 1, 1), (0, 1, 2)\}$, $V_3 = \{(0, 0, 1), (0, 0, 2)\}$ of $V(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3))$. V_1, V_2, V_3 are independent subsets and $V_1 \cup V_2 \cup V_3 = V(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3))$. We color the vertices in V_i with color i ($i = 1, 2, 3$). Therefore, the vertex chromatic number $\chi(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) \leq 3$. Since K_3 is subgraph, therefore, $\chi(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) \geq 3$. Hence, $\chi(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 3$.
- (iii) Since K_3 is subgraph, therefore, the clique number $\omega(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) \geq 3$. According to [6, Theorem 10.5], $\omega(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) \leq \chi(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 3$. Hence, $\omega(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 3$.
- (iv) Let $S_j \subset V(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3))$ be the set of elements with non-zero entry in j^{th} coordinate ($1 \leq j \leq 3$) and 0 or non-zero entries in the remaining coordinates. Then, $S_1 = \{(1, 0, 0), (1, 1, 0), (1, 0, 1), (1, 0, 2)\}$, $S_2 = \{(0, 1, 0), (1, 1, 0), (0, 1, 1), (0, 1, 2)\}$, $S_3 = \{(0, 0, 1), (1, 0, 1), (1, 0, 2), (0, 1, 1), (0, 1, 2)\}$ and $|S_1| = 4, |S_2| = 4, |S_3| = 6$. Therefore, S_3 is maximal independent subset and thus, the independence number, $\alpha(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 6 = \max \left\{ \frac{(|F_j|-1) \prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq 3 \right\}$.
- (v) Since, $\chi(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 3$, therefore, according to [11, Corollary 2.2], the metric chromatic number, $\mu(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 3$.
- (vi) Except the vertex $(1, 1, 0)$ which is of degree 2, all the other remaining vertices in $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ are odd degree vertices. Therefore, according to [9, Theorem 5.1], $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ is not Eulerian.

- (vii) $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ contains four vertices of degree 1. Therefore, there does not exist any cycle in $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ which contains all the vertices in $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$, since the degree 1 vertices can not belong to any cycle. In other words, $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ does not contain a Hamiltonian cycle. Therefore, $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ is not Hamiltonian.

Now, consider the complement graph $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$.




FIGURE 2. $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$

- (i) The clique number of $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$ is independence number of $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$ and independence number of $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$ is clique number of $\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)$, therefore, $\omega(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) = 6 = \max \left\{ \frac{(|F_j|-1) \prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq 3 \right\}$ and $\alpha(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) = 3$.
- (ii) Consider the partition $V_1 = \{(1, 0, 0), (0, 1, 0), (0, 0, 1)\}$, $V_2 = \{(0, 0, 2), (1, 1, 0)\}$, $V_3 = \{(0, 1, 1)\}$, $V_4 = \{(0, 1, 2)\}$, $V_5 = \{(1, 0, 1)\}$, $V_6 = \{(0, 1, 2)\}$. Then $V_1, V_2, V_3, V_4, V_5, V_6$ are independent subsets and $V_1 \cup V_2 \cup V_3 \cup V_4 \cup V_5 \cup V_6 = V(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)})$. We color vertices in V_i with color i ($1 \leq i \leq 6$). Therefore, $\chi(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) \leq 6$. Also, $\chi(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) \geq \omega(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) = 6$. Hence, $\chi(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) = 6$.
- (iii) In [13, Remark 1], Akbari and Mohammadian proved that if G is a graph such that for every vertex u of maximum degree there exists an edge uv such that $\Delta(G) - d(v) + 2$ is more than the number of vertices with maximum degree in G , then $\chi_1(G) = \Delta(G)$. In $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$, there are four vertices of maximum degree, namely, $(0, 1, 1), (0, 1, 2), (1, 0, 1), (1, 0, 2)$. Consider $(0, 1, 1)$. We have $\deg(0, 1, 1) = 7$. The vertex $(0, 1, 0)$ is adjacent to $(0, 1, 1)$ and $\deg(0, 1, 0) = 3$. Therefore, if $u = (0, 1, 1), v = (0, 1, 0)$, then $\Delta(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) - d(v) + 2 = 7 - 3 + 2 = 6$. Similarly for $u = (0, 1, 2)$ take $v = (0, 1, 0)$, for $u = (1, 0, 1)$

take $v = (1, 0, 0)$, for $u = (1, 0, 2)$ take $v = (1, 0, 0)$. Thus, according to [13, Remark 1], $\chi_1(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) = \Delta(\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}) = 7$. In other words, $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$ is Class-1.

- (iv) $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$ contains exactly one even degree vertex and all other vertices are odd degree vertices. Therefore, according to [7, Theorem 1.2.26], $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$ is not Eulerian. The cycle $(1, 0, 1), (1, 0, 0), (1, 1, 0), (0, 1, 0), (0, 1, 1), (0, 1, 2), (1, 0, 2), (0, 0, 1), (0, 0, 2), (1, 0, 1)$ is a Hamiltonian cycle in $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$. Therefore, $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)}$ is Hamiltonian.

2.1. Independence number, Clique number, Vertex Chromatic number. In Proposition 2.1, we find an upper bound for the independence number of $\Gamma(F_1 \times \cdots \times F_n)$. Note that if $n = 2$, then the independence number $\alpha(\Gamma(F_1 \times F_2)) = \max\{|F_1| - 1, |F_2| - 1\}$.

Proposition 2.1. *If $n \geq 3$, then the independence number $\alpha(\Gamma(F_1 \times \cdots \times F_n))$*

$$\geq \max\left\{\frac{(|F_j|-1)\prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq n\right\}.$$

Proof. First we observe that if two vertices are not adjacent in $\Gamma(F_1 \times \cdots \times F_n)$ they should have atleast one non-zero entry in the same coordinate, say j^{th} coordinate. Let S_j be the set of elements with non-zero entry in the j^{th} coordinate. Let $v = (a_1, \cdots, a_n) \in S_j$. The choices for a_j is $(|F_j| - 1)$ as $a_j \neq 0$ and choices for a_i is $|F_i|$ for $i \in \{1, \cdots, n\} \setminus \{j\}$ and S_j should contain elements of $V(\Gamma(F_1 \times \cdots \times F_n))$. Note that all coordinates having non-zero entries is not an element of $V(\Gamma(F_1 \times \cdots \times F_n))$ and we should remove the elements with each coordinate entry non-zero. The total number of choices of a_i with each of $a_i \neq 0$ is $(|F_i| - 1)$ and total such possible vertices not in $V(\Gamma(F_1 \times \cdots \times F_n))$ are $\prod_{i=1}^n (|F_i| - 1)$ and hence $|S_j| = \frac{(|F_j|-1)\prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1)$. Let $|S_\Delta| = \max\{|S_j| \mid 1 \leq j \leq n\}$ be set with largest cardinality. Hence, $\alpha(\Gamma(F_1 \times \cdots \times F_n)) \geq |S_\Delta| = \max\{|S_j| \mid 1 \leq j \leq n\} = \max\left\{\frac{(|F_j|-1)\prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq n\right\}$. $\square$

Note that if we consider $(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3))$, then

$$\alpha(\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_2 \times \mathbb{Z}_3)) = 6 = \max\left\{\frac{(|F_j|-1)\prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq 3\right\},$$

and if we consider $(\Gamma(\mathbb{Z}_5 \times \mathbb{Z}_5 \times \mathbb{Z}_5))$, then

$$\alpha(\Gamma(\mathbb{Z}_5 \times \mathbb{Z}_5 \times \mathbb{Z}_5)) = 48 > 36 = \max\left\{\frac{(|F_j|-1)\prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq 3\right\}.$$

In [1], Beck conjectured that the vertex chromatic number $\chi(R)$ is equal to clique number $\Omega(R)$ for a commutative ring R . Beck proved this conjecture in case of reduced ring [1, Theorem 3.8]. Beck proved that if R is a reduced ring $\neq (0)$ and if $\chi(R) < \infty$, then R has only a finite number of minimal prime ideals. And if n is this number then $\chi(R) = \Omega(R) = n + 1$. The result was proved using algebraic properties. Beck's result [1, Theorem 3.8] implies that chromatic number of the analogous zero divisor graph $\Gamma(R)$, as defined by Anderson and Livingston [2], is n .

In this paper, we provide graph theoretic approach for known results. We give the combinatorial proof for vertex chromatic number, clique number of $\Gamma(F_1 \times \cdots \times F_n)$ in

Proposition 2.2. Since the vertex set of $\Gamma(F_1 \times \cdots \times F_n)$ does not contain identity element $(0, \cdots, 0)$ and units of $F_1 \times \cdots \times F_n$, we use graph theoretical properties to prove $\chi(\Gamma(F_1 \times \cdots \times F_n)) = \omega(\Gamma(F_1 \times \cdots \times F_n)) = n$.

Proposition 2.2. (i) The vertex chromatic number $\chi(\Gamma(F_1 \times \cdots \times F_n)) = n$.

(ii) The clique number $\omega(\Gamma(F_1 \times \cdots \times F_n)) = n$.

Proof. (i) According to observation 2, $\Gamma(F_1 \times \cdots \times F_n)$ contains complete graph K_n as its subgraph. The vertex chromatic number $\chi(K_n)$ of the complete graph K_n is n . Therefore, the vertex chromatic number, $\chi(\Gamma(F_1 \times \cdots \times F_n)) \geq \chi(K_n) = n$. Now we partition $V(\Gamma(F_1 \times \cdots \times F_n))$ into n independent subsets. Let

$X_1 = \{v = (a_1, \cdots, a_n) \in V(\Gamma(F_1 \times \cdots \times F_n)) \mid a_1 \in F_1 \setminus \{0\}, a_j \in F_j, 2 \leq j \leq n\}$.

Then $|X_1| = (|F_1| - 1) \prod_{i=2}^n |F_i| - \prod_{i=1}^n (|F_i| - 1)$, since there are $(|F_1| - 1) \prod_{i=2}^n |F_i|$ number of possible choices for the n -tuple and among these n -tuple, $\prod_{i=1}^n (|F_i| - 1)$ are having non zero entries in each of n coordinate which are not elements of $V(\Gamma(F_1 \times \cdots \times F_n))$. Also the vertices in X_1 are not mutually adjacent to each other in $\Gamma(F_1 \times \cdots \times F_n)$.

Let $X_2 = \{v = (0, a_2, \cdots, a_n) \in V(\Gamma(F_1 \times \cdots \times F_n)) \mid a_2 \in F_2 \setminus \{0\}, a_j \in F_j, 3 \leq j \leq n\}$.

Then $|X_2| = (|F_2| - 1) \prod_{i=3}^n |F_i|$. Also, the vertices in X_2 are not mutually adjacent to each other in $\Gamma(F_1 \times \cdots \times F_n)$ and $X_1 \cap X_2 = \emptyset$. Similarly, let $X_k = \{v = (0, \cdots, 0, a_k, \cdots, a_n) \in V(\Gamma(F_1 \times \cdots \times F_n)) \mid a_k \in F_k \setminus \{0\}, a_j \in F_j, k+1 \leq j \leq n\}$.

Then $|X_k| = (|F_k| - 1) \prod_{i=k+1}^n |F_i|$ and the vertices in X_k are not mutually adjacent to each other in $\Gamma(F_1 \times \cdots \times F_n)$. Also, $X_i \cap X_j = \emptyset$ for $1 \leq i \neq j \leq k$. Therefore,

$$|X_1| + |X_2| + \cdots + |X_k| + \cdots + |X_{n-1}| + |X_n| \\ = [(|F_1| - 1) \prod_{i=2}^n |F_i| - \prod_{i=1}^n (|F_i| - 1)] + (|F_2| - 1) \prod_{i=3}^n |F_i| + \cdots + (|F_k| - 1) \prod_{i=k+1}^n |F_i| + \cdots + (|F_{n-1}| - 1) |F_n| + (|F_n| - 1)$$

$$= [\prod_{i=1}^n |F_i| - \prod_{i=2}^n |F_i|] + [\prod_{i=2}^n |F_i| - \prod_{i=3}^n |F_i|] + \cdots + [\prod_{i=k}^n |F_i| - \prod_{i=k+1}^n |F_i|] + \cdots + [|F_{n-1}| |F_n| - |F_n|] + (|F_n| - 1) - \prod_{i=1}^n (|F_i| - 1)$$

$$= \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1$$

$$= |V(\Gamma(F_1 \times F_2 \times \cdots \times F_n))|. \quad [10, \text{Theorem 2.1}]$$

Thus, $X_i, (1 \leq i \leq n)$ is a partition of $V(\Gamma(F_1 \times \cdots \times F_n))$ such that,

$\cup_{i=1}^n X_i = V(\Gamma(F_1 \times \cdots \times F_n))$ and moreover each $X_i, (1 \leq i \leq n)$ is an independent set. If we color the vertices in set $X_i, (1 \leq i \leq n)$ with color $i, (1 \leq i \leq n)$, we get a n -coloring of $\Gamma(F_1 \times \cdots \times F_n)$. Therefore, $\chi(\Gamma(F_1 \times \cdots \times F_n)) \leq n$. Since, we have already proved that $\chi(\Gamma(F_1 \times \cdots \times F_n)) \geq \chi(K_n) = n$, we have that the vertex chromatic number $\chi(\Gamma(F_1 \times \cdots \times F_n)) = n$.

(ii) Since, K_n is a subgraph of $\Gamma(F_1 \times \cdots \times F_n)$, therefore, $\omega(\Gamma(F_1 \times \cdots \times F_n)) \geq n$. According to [6, Theorem 10.5], $\omega(\Gamma(F_1 \times \cdots \times F_n)) \leq \chi(\Gamma(F_1 \times \cdots \times F_n)) = n$. Hence, $\omega(\Gamma(F_1 \times \cdots \times F_n)) = n$. $\square$

Corollary 2.1. (i) The vertex cover number, $\tau(\Gamma(F_1 \times \cdots \times F_n))$

$$\leq [\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1] - \max \left\{ \frac{(|F_j| - 1) \prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq n \right\}.$$

(ii) The clique number of complement graph,

$$\omega(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \geq \max \left\{ \frac{(|F_j| - 1) \prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq n \right\}.$$

- (iii) The independence number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is $\alpha(\overline{\Gamma(F_1 \times \cdots \times F_n)}) = n$.
- (iv) The vertex cover number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is

$$\tau(\overline{\Gamma(F_1 \times \cdots \times F_n)}) = [\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1] - n.$$
- (v) The vertex chromatic number of complement graph,

$$\chi(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \geq \max \left\{ (|F_j| - 1) \frac{\prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq n \right\}.$$

Proof. Using the fact, $|V(\Gamma(F_1 \times \cdots \times F_n))| = \alpha(\overline{\Gamma(F_1 \times \cdots \times F_n)}) + \tau(\overline{\Gamma(F_1 \times \cdots \times F_n)})$ [7], result (i) follows. Since the clique number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is independence number of $\Gamma(F_1 \times \cdots \times F_n)$ and independence number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is clique number of $\Gamma(F_1 \times \cdots \times F_n)$, and using the fact, $|V(\overline{\Gamma(F_1 \times \cdots \times F_n)})| = \alpha(\overline{\Gamma(F_1 \times \cdots \times F_n)}) + \tau(\overline{\Gamma(F_1 \times \cdots \times F_n)})$ [7], the result (ii), (iii), (iv) follows.

- (v) Follows from [6, Theorem 10.5] and (ii), $\chi(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \geq \omega(\overline{\Gamma(F_1 \times \cdots \times F_n)})$

$$\geq \max \left\{ \frac{(|F_j| - 1) \prod_{i=1}^n |F_i|}{|F_j|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq j \leq n \right\}. \quad \square$$

2.2. Metric Chromatic number.

Definition 2.1. For a set $S \subseteq V(G)$ and a vertex v of G , the distance $d(v, S)$ between v and S is defined as $d(v, S) = \min\{d(v, x) \mid x \in S\}$. [6, 9] Suppose that $c : V(G) \rightarrow \{1, 2, \dots, k\}$ is a k -coloring of G for some positive integer k where adjacent vertices may be colored the same and let $V_1, V_2, \dots, V_k$ be the resulting color classes. With each vertex v , we can associate a k -vector

$$\text{code}(v) = (a_1, a_2, \dots, a_k) = a_1 a_2 \cdots a_k$$

called the metric color code of v , where for each i with $1 \leq i \leq k$, $a_i = d(v, V_i)$. If $\text{code}(u) \neq \text{code}(v)$ for every two adjacent vertices u and v of G , then c is called a metric coloring of G . The minimum k for which G has a metric k -coloring is called the metric chromatic number of G and is denoted by $\mu(G)$. Clearly, $\mu(G)$ is defined for every connected graph G and $\mu(G) \geq 2$ for every nontrivial connected graph G . [11]

Let c be a proper k -coloring of a nontrivial connected graph G with resulting color classes $V_1, V_2, \dots, V_k$ and let u and v be two adjacent vertices of G . Then $u \in V_i$ and $v \in V_j$ for some $i, j \in \{1, 2, \dots, k\}$ with $i \neq j$. Suppose that $\text{code}(u) = (a_1, a_2, \dots, a_k)$ and $\text{code}(v) = (b_1, b_2, \dots, b_k)$. Then $a_i = b_j = 0$ and $a_j = b_i = 1$. Thus $\text{code}(u) \neq \text{code}(v)$ and c is also a metric coloring of G . Thus $\mu(G) \leq \chi(G)$. [11]

Lemma 2.1. If G be a connected graph on $n \geq 2$ vertices which contains complete graph K_p ; ($2 \leq p \leq n$) as an induced subgraph, then the metric chromatic number $\mu(G) \geq p$.

Proof. Suppose $\mu(G) = t$, $2 \leq t \leq p - 1$. Then there exists a metric t -coloring $c : V(G) \rightarrow \{1, 2, \dots, t\}$, such that, $\text{code}(u) \neq \text{code}(v)$ for every two adjacent vertices u and v of G . Then, $c|_{V(K_p)} : V(K_p) \rightarrow \{1, 2, \dots, t\}$ is also a t -metric coloring of induced subgraph K_p . But this is a contradiction, as a connected graph G of order n has metric chromatic number n if and only if $G = K_n$ [11]. Hence, $\mu(G) \geq p$. $\square$

Theorem 2.1. The metric chromatic number of $\Gamma(F_1 \times \cdots \times F_n)$ is $\mu(\Gamma(F_1 \times \cdots \times F_n)) = n$.

Proof. According to [11, Proposition 2.1], if $n = 2$, then $\mu(\Gamma(F_1 \times F_2)) = 2$, and if $n = 3$, then according to Proposition 2.2, $\chi(\Gamma(F_1 \times F_2 \times F_3)) = 3$, therefore, according to [11, Corollary 2.2], $\mu(\Gamma(F_1 \times F_2 \times F_3)) = 3$. Let $n \geq 4$. Claim: $\mu(\Gamma(F_1 \times \cdots \times F_n)) = n$. Suppose $\mu(\Gamma(F_1 \times \cdots \times F_n)) = t$, ($4 \leq t \leq n - 1$). Then there exists a metric t -coloring $c : V(\Gamma(F_1 \times \cdots \times F_n)) \rightarrow \{1, 2, \dots, t\}$ where adjacent vertices may be colored the same and

$code(u) \neq code(v)$ for every two adjacent vertices u and v of $\Gamma(F_1 \times \cdots \times F_n)$. According to observation 2, K_n is a subgraph of $\Gamma(F_1 \times \cdots \times F_n)$. Therefore, according to 2.1, $c|_{V(K_n)} : V(K_n) \rightarrow \{1, 2, \dots, t\}$ is a metric t -coloring of the subgraph K_n , such that for every pair x, y of adjacent vertices in subgraph K_n , $code(x) \neq code(y)$. This is a contradiction to the fact that a connected graph G of order n has metric chromatic number n if and only if $G = K_n$ [11]. Thus, $\mu(\Gamma(F_1 \times \cdots \times F_n)) \geq n$. According to [11], since metric chromatic number $\leq$ vertex chromatic number, we have, $\mu(\Gamma(F_1 \times \cdots \times F_n)) \leq \chi(\Gamma(F_1 \times \cdots \times F_n)) = n$. Hence, $\mu(\Gamma(F_1 \times \cdots \times F_n)) = \chi(\Gamma(F_1 \times \cdots \times F_n)) = n$. $\square$

Corollary 2.2. *The metric chromatic number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is $\mu(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \geq \max \left\{ (|F_k| - 1) \frac{\prod_{i=1}^n |F_i|}{|F_k|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq k \leq n \right\}$.*

Proof. Let $\omega(\overline{\Gamma(F_1 \times \cdots \times F_n)}) = t$. According to 2.1, $\mu(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \geq t$. Therefore, according to Corollary 2.1 (ii), $\mu(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \geq t$
 $\geq \max \left\{ (|F_k| - 1) \frac{\prod_{i=1}^n |F_i|}{|F_k|} - \prod_{i=1}^n (|F_i| - 1) \mid 1 \leq k \leq n \right\}$. $\square$

2.3. Edge Chromatic number. In [12], Akbari and Mohammadian, have shown that for any finite commutative ring R , the edge chromatic number of $\Gamma(R)$ is equal to the maximum degree of $\Gamma(R)$, unless $\Gamma(R)$ is a complete graph of odd order. They have also proved in [12, Remark 1], that if G is a graph such that for every vertex u of maximum degree there exists an edge uv such that $\Delta(G) - d(v) + 2$ is more than the number of vertices with maximum degree in G , then $\chi_1(G) = \Delta(G)$. We use this remark to determine the edge chromatic number of $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. Note that if $n = 2$, then $\overline{\Gamma(F_1 \times F_2)}$ is disjoint union of two complete graphs $K_{|F_1|-1}$ and $K_{|F_2|-1}$. Thus, $\overline{\Gamma(F_1 \times F_2)}$ is Class-1 if $\max\{|F_1| - 1, |F_2| - 1\}$ is even, and Class-2 if $\max\{|F_1| - 1, |F_2| - 1\}$ is an odd number greater than or equal to 3. [6, Theorem 10.15]

Theorem 2.2. $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ belongs to Class-1

- (i) if $n \geq 3$ and all the fields are of order two
- (ii) if there are at least two fields of order two and at least one field of order greater than two.

Proof. (i) If all the fields are of order two, then $\delta(\Gamma(F_1 \times \cdots \times F_n)) = 1$, therefore, $\Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)}) = |V(\Gamma(F_1 \times \cdots \times F_n))| - 1 - 1 = (2^n - 2) - 2 = 2^n - 4$. If u is a vertex with exactly one 0 in i^{th} ($1 \leq i \leq n$) coordinate and 1^s in the remaining coordinates, then $deg_{\Gamma(F_1 \times \cdots \times F_n)}(u) = 1$ and thus, $deg_{\overline{\Gamma(F_1 \times \cdots \times F_n)}}(u) = 2^n - 4 = \Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)})$. It follows that if order of each field is two, then there are exactly n maximum degree vertices in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. Let v be a vertex with exactly one 1 in j^{th} ($1 \leq j \neq i \leq n$) coordinate and 0^s in the remaining coordinates, then $deg_{\Gamma(F_1 \times \cdots \times F_n)}(v) = 2^{n-1} - 1$ and thus, $deg_{\overline{\Gamma(F_1 \times \cdots \times F_n)}}(v) = (2^n - 2) - (2^{n-1} - 1) - 1 = 2^{n-1} - 2$. Therefore, $\Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)}) - deg(v) + 2 = (2^n - 4) - (2^{n-1} - 2) + 2 = 2^{n-1} > n$ if $n \geq 3$. Thus, it follows that, if all the fields are of order two, then $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ belongs to Class-1.

(ii) Suppose there are at least two fields of order two and at least one field of order greater than two. Suppose there are t ($2 \leq t \leq n - 1$) fields of order two. Let $F_{j_1}, \dots, F_{j_t}$ ($1 \leq t \leq n - 1$) be the t fields of order two. Let u be a vertex with 0 in exactly one coordinate j_s ($1 \leq s \leq t$) and non-zero entries in the remaining coordinates. Then $deg_{\Gamma(F_1 \times \cdots \times F_n)}(u) = |F_{j_s}| - 1 = 2 - 1 = 1$. Therefore, $deg_{\overline{\Gamma(F_1 \times \cdots \times F_n)}}(u) =$

$|V(\Gamma(F_1 \times \cdots \times F_n))| - \deg_{\Gamma(F_1 \times \cdots \times F_n)}(u) - 1$
 $= [\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1] - 1 - 1$
 $= \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 3 = \Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)}).$
 Let H_{j_s} ($1 \leq s \leq t$) be the set of vertices with exactly one 0 in the j_s coordinate and non-zero entries in the remaining $(n-1)$ coordinates. Then H_{j_s} contains maximum degree vertices and $|H_{j_s}| = \prod_{i=1, i \neq j_s}^n (|F_i| - 1)$. Note that $H_{j_1} \cup \cdots \cup H_{j_t}$ is the set containing maximum degree vertices and $|H_{j_1} \cup \cdots \cup H_{j_t}|$
 $= \prod_{i=1, i \neq j_1}^n (|F_i| - 1) + \cdots + \prod_{i=1, i \neq j_t}^n (|F_i| - 1)$
 $= \prod_{i=1, i \neq j_1, \dots, j_t}^n (|F_i| - 1) + \cdots + \prod_{i=1, i \neq j_1, \dots, j_t}^n (|F_i| - 1)$
 $= t \cdot \prod_{i=1, i \neq j_1, \dots, j_t}^n (|F_i| - 1).$
 Consider the set H_{j_s} . Let k be such that $|F_k| = 2$, $k \neq j_s$. Let v be a vertex with exactly one non-zero entry in the k^{th} coordinate and 0^s in the remaining coordinates. Since, the k^{th} coordinate of v and $u \in H_{j_s}$ is non-zero entry, therefore, $uv \in E(\overline{\Gamma(F_1 \times \cdots \times F_n)})$ and, $\deg_{\Gamma(F_1 \times \cdots \times F_n)}(v) = \prod_{i=1, i \neq k}^n |F_i| - 1$, therefore, $\deg_{\overline{\Gamma(F_1 \times \cdots \times F_n)}}(v)$
 $= |V(\Gamma(F_1 \times \cdots \times F_n))| - \deg_{\Gamma(F_1 \times \cdots \times F_n)}(v) - 1$
 $= [\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1] - [\prod_{i=1, i \neq k}^n |F_i| - 1] - 1$
 $= \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - \prod_{i=1, i \neq k}^n |F_i| - 1.$
 Therefore, $\Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)}) - \deg_{\overline{\Gamma(F_1 \times \cdots \times F_n)}}(v) + 2$
 $= [\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 3] - [\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - \prod_{i=1, i \neq k}^n |F_i| - 1] + 2$
 $= \prod_{i=1, i \neq k}^n |F_i|$
 $= 2^{t-1} \prod_{i=1, i \neq j_1, \dots, j_t}^n |F_i|$ (since $k \in \{j_1, \dots, j_t\} \setminus j_s$)
 $> t \prod_{i=1, i \neq j_1, \dots, j_t}^n (|F_i| - 1)$
 (since $2^{t-1} > t$ if $t \geq 3$ and for $t = 2$, $|H_{j_1} \cup H_{j_2}| = 2 \prod_{i=1, i \neq j_1, j_2}^n (|F_i| - 1)$ and there exists at least one field of order greater than two.)
 Thus, according to [12, Remark 1], $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ belongs to Class-1, if there are at least two fields of order two and at least one field of order greater than two. $\square$

Remark 2.1. (i) The technique mentioned in the above proof ([12, Remark 1]) holds for $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_3 \times \mathbb{Z}_3)}$ but does not hold for $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_5 \times \mathbb{Z}_5)}$, $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_7 \times \mathbb{Z}_7)}$, $\overline{\Gamma(\mathbb{Z}_2 \times \mathbb{Z}_{11} \times \mathbb{Z}_{11})}$, and so on.

(ii) Suppose each field is of order p , where p is an odd prime number. Consider $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. Then $\delta(\Gamma(F_1 \times \cdots \times F_n)) = p-1$, therefore, $\Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)}) = |V(\Gamma(F_1 \times \cdots \times F_n))| - (p-1) - 1 = (p^n - (p-1)^n - 1) - (p-1) - 1 = p^n - (p-1)^n - p - 1$. Let H_j ($1 \leq j \leq n$) be the set of vertices with exactly one 0 in the j^{th} coordinate and non-zero entries in the remaining $(n-1)$ coordinates. Then $H_1 \cup \cdots \cup H_n$ is the set containing maximum degree vertices in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ and $|H_1 \cup \cdots \cup H_n| = n \cdot (p-1)^{n-1}$. Consider the set H_j . Let v be a vertex with exactly one non-zero entry in k^{th} ($1 \leq k \neq j \leq n$) coordinate and 0^s in the remaining coordinates, then $\deg_{\Gamma(F_1 \times \cdots \times F_n)}(v) = p^{n-1} - 1$ and thus, $\deg_{\overline{\Gamma(F_1 \times \cdots \times F_n)}}(v)$
 $= (p^n - (p-1)^n - 1) - (p^{n-1} - 1) - 1 = p^n - (p-1)^n - p^{n-1} - 1$. Therefore,
 $\Delta(\overline{\Gamma(F_1 \times \cdots \times F_n)}) - \deg(v) + 2$
 $= (p^n - (p-1)^n - p - 1) - (p^n - (p-1)^n - p^{n-1} - 1) + 2 = p^{n-1} - p + 2.$
 Thus, it follows that $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ belongs to Class-1 if

$$p^{n-1} - p + 2 > n \cdot (p-1)^{n-1}.$$

We observe that for a fixed prime p , the above inequality holds for large n . (For example, if $p = 3$ and $n \geq 6$, if $p = 5$ and $n \geq 13$, if $p = 7$ and $n \geq 21$, if $p = 11$ and $n \geq 40$,

and so on.) For a fixed prime p , the term p^{n-1} in $p^{n-1} - p + 2$ grows exponentially much faster than both the linear and exponential factors in $n(p-1)^{n-1}$, as $n \rightarrow \infty$. This means that for large n , the left-hand side of the inequality becomes larger than the right-hand side. Therefore, for each prime p , there exists a natural number $n_0(p)$, such that, for all $n \geq n_0(p)$, the inequality $p^{n-1} - p + 2 > n \cdot (p-1)^{n-1}$ is always true for all $n \geq n_0(p)$. In other words, if all the fields are of order p , where $p \geq 3$ is a prime number, then there exists a natural number $n_0(p)$, such that, for all $n \geq n_0(p)$, $\Gamma(F_1 \times \cdots \times F_n)$ belongs to Class-1. If $|F_1| \leq |F_2| \leq \cdots \leq |F_n|$ and $|F_1| = p_1$ is an odd prime, then there exists a natural number $n_0(p_1)$, such that, for all $n \geq n_0(p_1)$, $\Gamma(F_1 \times \cdots \times F_n)$ belongs to Class-1.

2.4. Eulerian and Hamiltonian. In [13], S. Akbari, A. Mohammadian, defined the zero-divisor graph of a ring R as the directed graph $\Gamma(R)$, such that its vertices are all non-zero zero-divisors of R in which for any two distinct vertices x and y , $x \rightarrow y$ is an edge if and only if $xy = 0$. They proved that [13, Proposition 1], if $n \geq 2$ and $R_1, \dots, R_n$ are finite rings and $R = R_1 \times \cdots \times R_n$, then $\Gamma(R)$ is an Eulerian graph if and only if for $i = 1, \dots, n$, either R_i is a field or $\Gamma(R_i)$ is an Eulerian graph. In this study, the zero divisor graph (defined by Anderson and Livingston [2]) we are considering is a simple undirected graph. In Proposition 2.3, we determine the conditions for which the zero divisor graph $\Gamma(F_1 \times \cdots \times F_n)$ is Eulerian or not Eulerian. The conditions are determined in terms of the order of the fields. Since the order of a finite field is a prime power, the following proposition fully characterizes the Eulerian property for the graphs under consideration in this study.

Proposition 2.3. *If $n \geq 2$ then $\Gamma(F_1 \times \cdots \times F_n)$ is*

- (i) *not Eulerian if $|F_i| = 2^r$, for some i , $(1 \leq i \leq n)$, $r \in \mathbb{N}$.*
- (ii) *Eulerian if $|F_i| \neq 2^r$ for all i , $(1 \leq i \leq n)$.*

Proof. Let $v \in V(\Gamma(F_1 \times \cdots \times F_n))$. Suppose v contains 0^s in coordinate $i_1, i_2, \dots, i_r$ with $1 \leq i_1 < i_2 < \cdots < i_r \leq n$. Then $\deg(v) = \prod_{k=1}^r |F_{i_k}| - 1$.

(i) If $|F_i| = 2^r$, for some i , $(1 \leq i \leq n)$, $r \in \mathbb{N}$, then clearly the degree of a vertex with 0 in i^{th} coordinate and non-zero entries in remaining coordinate is odd and thus $\Gamma(F_1 \times \cdots \times F_n)$ is not Eulerian.[9]

(ii) If $|F_i| \neq 2^r$ for all i , $(1 \leq i \leq n)$ then every vertex in $\Gamma(F_1 \times \cdots \times F_n)$ is of even degree. Hence according to [7, Theorem 1.2.26], $\Gamma(F_1 \times \cdots \times F_n)$ is Eulerian in this case. $\square$

In Proposition 2.4, we determine the conditions for $\Gamma(F_1 \times \cdots \times F_n)$ to be Hamiltonian or non-Hamiltonian. We make use of the result [7, Proposition 7.2.3], "If a graph G has a Hamiltonian cycle then for each non-empty subset $S \subseteq V(G)$ the graph $G - S$ has at most $|S|$ components. "

Proposition 2.4. (i) $\Gamma(F_1 \times F_2)$ is Hamiltonian if $|F_1| = |F_2| \geq 3$, and $\Gamma(F_1 \times F_2)$ is not Hamiltonian if $|F_2| > |F_1|$ or $|F_1| > |F_2|$.
(ii) $\Gamma(F_1 \times \cdots \times F_n)$ is not Hamiltonian, if $n \geq 3$.

Proof. (i) $\Gamma(F_1 \times F_2)$ is complete bipartite graph $K_{|F_1|-1, |F_2|-1}$. A complete bipartite graph is hamiltonian if and only if the parts are of equal size and have 2 or more vertices[7]. Therefore, $K_{|F_1|-1, |F_2|-1}$ is Hamiltonian if and only if $|F_1| - 1 = |F_2| - 1 \geq 2$. Hence, the result follows.

(ii) Let $n \geq 3$. Without loss of generality, let $|F_1| = \min\{|F_1|, \dots, |F_n|\}$.

Consider the set $S = \{(t, 0, \dots, 0) \mid t \in F_1 \setminus \{0\}\}$. We have $|S| = |F_1| - 1$.

Let $T = \{(0, a_2, \dots, a_n) : a_j \in \{1, 2, \dots, |F_j| - 1\}, (2 \leq j \leq n)\}$.

Then, $|T| = \prod_{j=2}^n (|F_j| - 1)$. Each vertex in S is adjacent with each and every vertex in T and the only vertices adjacent to a vertex of T are in S . That is, the vertices in T are not adjacent with any vertex in $\Gamma(F_1 \times \cdots \times F_n) - S \implies \deg(x) = |F_1| - 1$ for $x \in T$. The number of components in $\Gamma(F_1 \times \cdots \times F_n) - S \geq \prod_{j=2}^n (|F_j| - 1) > |F_1| - 1 = |S|$, as $|F_1| = \min\{|F_1|, \dots, |F_n|\}$. Hence, if $n \geq 3$, then according to [7, Proposition 7.2.3], $\Gamma(F_1 \times \cdots \times F_n)$ is not Hamiltonian. $\square$

In Proposition 2.5, we determine the conditions for $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ to be Eulerian, Hamiltonian. In [14, Corollary 3.5], Devhare, Joshi, Lagrange gave the conditions for $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ to be Eulerian. We prove the same result using graph theoretic approach.

Proposition 2.5. (i) $\overline{\Gamma(F_1 \times F_2)}$ is neither Eulerian nor Hamiltonian.
(ii) $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is Hamiltonian if $n \geq 3$.
(iii) $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is Eulerian if and only if $|F_i| = 2^{s_i}$, $s_i \geq 1$, $\forall 1 \leq i \leq n$, $n \geq 3$.

Proof. (i) $\overline{\Gamma(F_1 \times F_2)}$ is disconnected.

(ii) Let $n \geq 3$. We prove that $\overline{\Gamma(F_1 \times \cdots \times F_n)}$ is Hamiltonian. We demonstrate a Hamiltonian cycle in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. First we partition $V(\overline{\Gamma(F_1 \times \cdots \times F_n)})$ into n subsets such that each subset is a clique in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$.

Let $X_1 = \{(a_1, a_2, \dots, a_n) \in V(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \mid a_1 \in F_1 \setminus \{0\}, a_j \in F_j, 2 \leq j \leq n\}$. Then $|X_1| = (|F_1| - 1) \prod_{i=2}^n |F_i| - \prod_{i=1}^n (|F_i| - 1)$, since there are $(|F_1| - 1) \prod_{i=2}^n |F_i|$ number of possible choices for the n -tuple and among these n -tuple, $\prod_{i=1}^n (|F_i| - 1)$ are having non zero entries in each of n coordinate which are not elements of $V(\overline{\Gamma(F_1 \times \cdots \times F_n)})$. Also the vertices in X_1 are all mutually adjacent to each other in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$.

Let $X_2 = \{(0, a_2, \dots, a_n) \in V(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \mid a_2 \in F_2 \setminus \{0\}, a_j \in F_j, 3 \leq j \leq n\}$. Then $|X_2| = (|F_2| - 1) \prod_{i=3}^n |F_i|$. And the vertices in X_2 are all mutually adjacent to each other in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. Also, $X_1 \cap X_2 = \emptyset$. Similarly, let $X_k = \{(0, \dots, 0, a_k, \dots, a_n) \in V(\overline{\Gamma(F_1 \times \cdots \times F_n)}) \mid a_k \in F_k \setminus \{0\}, a_j \in F_j, k+1 \leq j \leq n\}$.

Then $|X_k| = (|F_k| - 1) \prod_{i=k+1}^n |F_i|$. And the vertices in X_k are all mutually adjacent to each other in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$. Also, $X_i \cap X_j = \emptyset$ for $1 \leq i \neq j \leq k$. Therefore,
 $|X_1| + |X_2| + \cdots + |X_k| + \cdots + |X_{n-1}| + |X_n|$

$$= \left[(|F_1| - 1) \prod_{i=2}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) \right] + (|F_2| - 1) \prod_{i=3}^n |F_i| + \cdots + (|F_k| - 1) \prod_{i=k+1}^n |F_i| + \cdots + (|F_{n-1}| - 1) |F_n| + (|F_n| - 1)$$

$$= \left[\prod_{i=1}^n |F_i| - \prod_{i=2}^n |F_i| \right] + \left[\prod_{i=2}^n |F_i| - \prod_{i=3}^n |F_i| \right] + \cdots + \left[\prod_{i=k}^n |F_i| - \prod_{i=k+1}^n |F_i| \right] + \cdots + \left[|F_{n-1}| |F_n| - |F_n| \right] + (|F_n| - 1) - \prod_{i=1}^n (|F_i| - 1)$$

$$= \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1$$

$$= |V(\overline{\Gamma(F_1 \times \cdots \times F_n)})|.$$

Thus, X_i , $(1 \leq i \leq n)$ is a partition of $V(\overline{\Gamma(F_1 \times \cdots \times F_n)})$ such that,

$\cup_{i=1}^n X_i = V(\overline{\Gamma(F_1 \times \cdots \times F_n)})$ and moreover each X_i , $(1 \leq i \leq n)$ is a clique set in $\overline{\Gamma(F_1 \times \cdots \times F_n)}$.

Let P_1 be a Hamiltonian path of vertices in X_1 such that the starting vertex of path P_1 contains 1 in the first coordinate and last n^{th} coordinate and the last vertex of path P_1 contains 1 in the second coordinate. Let P_2 be a Hamiltonian path of vertices in X_2

such that the last vertex of path P_2 contains 1 in the third coordinate. Clearly, the last vertex of path P_1 is adjacent to first vertex of path P_2 . Therefore, $P_1 \cup P_2$ is a Hamiltonian path of vertices in $X_1 \cup X_2$. Similarly, let P_k be a Hamiltonian path of vertices in X_k ($3 \leq k \leq n-1$) such that the last vertex of path P_k contains 1 in the $k+1$ coordinate. Clearly, the last vertex of path P_k is adjacent to first vertex of path P_{k+1} and $P_1 \cup P_2 \cup \dots \cup P_k$ is a Hamiltonian path of vertices in $X_1 \cup X_2 \cup \dots \cup X_k$. Let P_n be a Hamiltonian path of vertices in X_n . The vertices of X_n contain non-zero entry in the last n^{th} coordinate. Therefore, the last vertex of path P_n is adjacent to the first vertex of path P_1 . Therefore, $P_1 \cup P_2 \cup \dots \cup P_n$ is a Hamiltonian path of vertices in $V(\Gamma(F_1 \times \dots \times F_n))$ such that the last vertex of path P_n is adjacent to first vertex of path P_1 . Thus, we get a Hamiltonian cycle in $\overline{\Gamma(F_1 \times \dots \times F_n)}$. Hence, $\overline{\Gamma(F_1 \times \dots \times F_n)}$ is Hamiltonian.

(iii) Let $n \geq 3$. Let $x \in V(\Gamma(F_1 \times \dots \times F_n))$ contains 0^s in the coordinates $i_1 \dots, i_r$. Then $\deg_{\Gamma(F_1 \times \dots \times F_n)}(x) = \prod_{i_k=1}^r |F_{i_k}| - 1$ and $|V(\Gamma(F_1 \times \dots \times F_n))| = |V(\overline{\Gamma(F_1 \times \dots \times F_n)})| = \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - 1$. Hence $\deg_{\overline{\Gamma(F_1 \times \dots \times F_n)}}(x) = \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - \prod_{k=1}^r |F_{i_k}| - 1$ where $1 \leq i_1 < i_2 < \dots < i_r \leq n$, ($1 \leq r \leq n-1$). If $|F_i| = 2^{s_i}$, $s_i \geq 1$ for all $1 \leq i \leq n$, then, $\prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - \prod_{k=1}^r |F_{i_k}| - 1$ is even, that is, the degree of every vertex is even and hence $\overline{\Gamma(F_1 \times \dots \times F_n)}$ is Eulerian in this case.

Suppose at least one field is of odd order. There are two possibilities:

- (a) All the fields are of odd order.
- (b) At least one field is of odd order and at least one field of even order.

(a) If all the fields are of odd order, that is, $|F_j| = p_j^{s_j}$, p_j - prime, $p_j \neq 2$, $s_j \geq 1$ for all $1 \leq j \leq n$, then for $x \in V(\Gamma(F_1 \times \dots \times F_n))$ containing 0^s in the coordinates $i_1 \dots, i_r$, $\deg_{\overline{\Gamma(F_1 \times \dots \times F_n)}}(x) = \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - \prod_{k=1}^r |F_{i_k}| - 1$ is odd.

(b) If $|F_j| = p_j^{s_j}$, p_j - prime, $p_j \neq 2$, $s_j \geq 1$ for some $1 \leq j \leq n$ and $|F_k| = 2^{s_k}$, $s_k \geq 1$ for some $1 \leq k \neq j \leq n$, then let y be a vertex with 0 in k^{th} coordinate and non-zero entries in the remaining coordinates. Then $\deg_{\overline{\Gamma(F_1 \times \dots \times F_n)}}(y) = \prod_{i=1}^n |F_i| - \prod_{i=1}^n (|F_i| - 1) - |F_k| - 1$ is odd. Thus, it follows that, $\overline{\Gamma(F_1 \times \dots \times F_n)}$ is not Eulerian if at least one field is of odd order. $\square$

CONCLUSION

We obtained an upper bound for the independence number and determined the Eulerian and Hamiltonian properties of the zero-divisor graph $\Gamma(F_1 \times \dots \times F_n)$ and the complement graph $\overline{\Gamma(F_1 \times \dots \times F_n)}$. We determined the Edge chromatic number of $\overline{\Gamma(F_1 \times \dots \times F_n)}$ and proved that the metric chromatic number clique number and vertex chromatic number of $\Gamma(F_1 \times \dots \times F_n)$ is n and also determined clique number, vertex chromatic number, metric chromatic number of $\overline{\Gamma(F_1 \times \dots \times F_n)}$.

Acknowledgement: The authors would like to extend their gratitude to the anonymous referees for their valuable suggestions.

Declarations :

- Declarations of interest: None.
- Conflict of interest: On behalf of all authors, the corresponding author states that there is no conflict of interest.

- Authors' contributions. Dr. Nithya Sai Narayana and Subhash Mallinath Gaded wrote the main manuscript text and reviewed the manuscript. Both the authors contributed equally to these work. Both authors have read and approved the final manuscript.

Data Availability : No data was used for conduct of this study.

Funding : No funding was received for conducting this study.

Declaration of Generative AI and AI-assisted technologies in the writing process :

The authors declare that they have not used the generative AI and AI-assisted technologies in the writing process.

REFERENCES

- [1] Beck, I., (1988), Coloring of commutative rings, *Journal of algebra*, 116(1), 208–226.
- [2] Anderson, D. F. and Livingston, P. S., (1999), The zero-divisor graph of a commutative ring, *Journal of Algebra*, 217(2), 434–447.
- [3] Atiyah, M., *Introduction To Commutative Algebra*, Taylor & Francis eBooks DRM Free Collection.
- [4] Gallian, J., (2021), *Contemporary abstract algebra*, Chapman and Hall/CRC.
- [5] Balakrishnan, R. and Ranganathan K., (2002), *A textbook of graph theory*, Springer Science & Business Media.
- [6] Chartrand, G. and Zhang, P., (2013), *A first course in graph theory*, Courier Corporation.
- [7] West, D. Brent and others, (2001(2)), *Introduction to graph theory*, Prentice hall Upper Saddle River.
- [8] Akbari, S., Ghanbari, M., Nikmehr, M. J., (2014), The chromatic index of a graph whose core is a cycle of order at most 13, *Graphs and Combinatorics*, 30(4), pp. 801–819.
- [9] Choudum, S. A., (1987), *A First Course in Graph Theory*, MacMillan India Limited.
- [10] Subhash, M. G., Nithya, S. N., (2023), On zero divisors graphs of direct product of finite fields, *Journal of Computational Mathematics*, 7(1), pp. 100–109.
- [11] Chartrand, G. and Okamoto, F. and Zhang, P., (2009), The metric chromatic number of a graph, *Australas. J Comb.*, 44, pp. 273–286.
- [12] Akbari, S and Mohammadian, A, (2004), On the zero-divisor graph of a commutative ring, *Journal of Algebra*, 274(2), pp. 847–855.
- [13] Akbari, S and Mohammadian, A, (2007), On zero-divisor graphs of finite rings, *Journal of Algebra*, 314(1), pp. 168–184.
- [14] Devhare, Sarika and Joshi, Vinayak and LaGrange, John D, (2019), Eulerian and Hamiltonian complements of zero-divisor graphs of pseudocomplemented posets, *Palestine Journal of Mathematics*, 8(2), pp. 30–39.



Dr. Nithya Sai Narayana is Associate Professor(Retd.) from N.E.S. Ratnam College of Arts Science and Commerce, Bhanudp(W), Mumbai, Maharashtra, India. She holds M.Phil and Ph.D. degree from Mumbai University. Her current research interests includes Extremal problems, Algebra, Algebraic graph theory, Crossing Numbers, Energy of graphs.



Subhash Mallinath Gaded received his M. Sc. and M.Phil. degree from University of Mumbai, Mumbai, Maharashtra, India. His research interests are Algebraic graph theory, zero divisor graphs, distances in graphs and topological indices. He is currently working as an Assistant professor in the Department of Mathematics, R K Talreja College of Arts, Science and Commerce, Ulhasnagar-03, District Thane, Maharashtra, India.